



IB 99
00844
02

The
Patent
Office

PCT / IB 99 / 0 0 8 4 4



31.05.99

INVESTOR IN PEOPLE

ESU

PRIORITY DOCUMENT

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP9 1RH

REC'D 02 JUN 1999


WIPO PCT

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.


Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.



Signed

P. Mahoney

Dated 10 May 1999



**The
Patent
Office**

Request for grant of a patent

(See notes on the back of this form. You can also get a explanatory leaflet from the Patent Office to help you fill in this form)

The Patent Office

Cardiff Road
Newport
Gwent NP9 1RH

1. Your reference **P19092GB G22187P**

2. Patent application number
(The Patent Office)

9808932.9

27 APR 1998

3. Full name, address and postcode of the or of each applicant (underline all surnames)

**CHIRON S.P.A.
Via Fiorentina 1
53100 Siena
ITALY**

Patents ADP number (if you know it)

If the applicant is a corporate body, give the country/state of its incorporation

ITALY

7157811001

4. Title of the invention **POLYEPITOPE CARRIER PROTEIN**

5. Name of your agent (if you have one)

Carpmaels & Ransford

"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)

**43 Bloomsbury Square
London
WC1A 2RA**

Patents ADP number (if you know it)

83001

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number

Country

Priority application number
(if you know it)

Date of filing
(day / month / year)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application

Number of earlier application

Date of filing
(day / month / year)

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer 'Yes' if:

- a) any applicant named in part 3 is not an inventor, or
- b) there is an inventor who is not named as an applicant, or
- c) any named applicant is a corporate body

Yes

See note (d))

Patents Form 1/77

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form

Description	54
Claim(s)	4
Abstract	
Drawing(s)	8



10. If you are also filing any of the following, state how many against each item.

Priority documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (*Patents Form 7/77*)

Request for preliminary examination and search (*Patents Form 9/77*)

Request for substantive examination (*Patents Form 10/77*)

Any other documents
(please specify)

11. I/We request the grant of a patent on the basis of this application.

Signature

Date

 27th April 1998

12. Name and daytime telephone number of person to contact in the United Kingdom
- | | |
|----------------------|---------------|
| HUW G. HALLYBONE | 0171 242 8692 |
| Carpmaels & Ransford | |

Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

Notes

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 0645 500505.
- Write your answers in capital letters using black ink or you may type them.
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- If you have answered 'Yes' Patents Form 7/77 will need to be filed.
- Once you have filled in the form you must remember to sign and date it.
- For details of the fee and ways to pay please contact the Patent Office.

Polyepitope Carrier Protein

The present invention relates to polyepitope carrier proteins. When conjugated to
5 capsular polysaccharides, these carrier proteins are useful as components of vaccines
that are capable of eliciting a T-cell dependent immune response. Particularly, the
proteins of the present invention may be used to confer protection against infection from
encapsulated bacteria in infants between the ages of 3 months and about 2 years.

10 Encapsulated bacteria such as *Haemophilus influenzae*, *Neisseria meningitidis* and
Streptococcus pneumoniae constitute a significant cause of morbidity and mortality in
neonates and infants world-wide (Tunkel & Scheld, 1993). In developing countries,
around one million children die each year due to pneumonia alone. Furthermore, even
in developed countries, the increase in the phenomenon of antibiotic resistance means
15 that there is a great need to improve on existing vaccines.

The polysaccharide capsule of *H. influenzae*, *N. meningitidis* and *S. pneumoniae*
represents a major virulence factor that is important for nasopharyngeal colonisation
and systemic invasion by encapsulated bacteria (Moxon and Kroll, 1990).
20 Consequently, much of the research directed towards finding protective immunogens
has focused on capsular polysaccharides. The finding that these polysaccharides are
able to elicit the formation of protective antibodies led to the development of a number
of vaccines that have been efficacious in protecting adults from disease (Andreoni *et al.*
1993; Goldblatt *et al.* 1992).

25

The problem with capsular polysaccharide vaccines developed to date is that they suffer
an inherent inability to protect children under two years of age from disease (Holmes
and Granoff 1992). This is a significant drawback when it is appreciated that this
population of children is at highest risk of infection. Their failure to block infection is
30 believed to derive from the T-cell independent (TI) type of immune reaction that is the
only antibody response used by the body against polysaccharide antigens. This type of
response does not involve MHC Class II restriction molecules for antigen presentation
to T-cells; as a consequence, T-cell help is prevented. Although the TI response works

well in adults, it is inactive in very young children due to a combination of factors such as functional B-cell immaturity, inactivation of B-cell receptor-mediated signalling and B-cell anergy in response to antigen stimulation.

- 5 To overcome this drawback, two particular vaccine approaches are currently being investigated. The first is the development of anti-idiotypic vaccines that contain peptides that mimic carbohydrate idiotypes (McNamara 1984; Agadjanyan, 1997). The second approach involves conjugate vaccines that are designed to transform T-cell independent (TI) polysaccharide antigens into T-dependent (TD) antigens through the
10 covalent linkage of the polysaccharide to a peptide carrier.

H. influenzae type B (Hib) conjugate vaccines represent a leading example for the development of other vaccines against infections that are due to capsulated bacteria. In fact, meningitis and other infections caused by Hib have declined dramatically in
15 countries where widespread vaccination with Hib conjugate has been achieved (Robins, 1996). Complete elimination of the pathogen might be possible, but depends upon several factors, including a further improvement of the existing vaccines (Liptak, 1997).

The widely distributed paediatric vaccine antigens tetanus and diphtheria toxoids have
20 been selected as carrier proteins with the aim of taking advantage of an already-primed population at the time of conjugate vaccine injection. Previous vaccination with paediatric diphtheria-tetanus (DT) or diphtheria-tetanus-pertussis (DTP) vaccines means that carrier priming may now be exploited to enhance the immune response to polysaccharide conjugates.

25 A number of such vaccines have been successfully produced and have been efficacious in reducing the number of deaths caused by these pathogens. The carriers used in these vaccines are large antigens such as tetanus toxoid, non-toxic diphtheria toxin mutant CRM197 and group B *N. meningitidis* outer membrane protein complex (OMPC).

- 30 However, in the future, it is thought that as the number of conjugate vaccines containing the same carrier proteins increases, the suppression of immune responses by pre-existing antibodies to the carrier is likely to become a problem.

Much research is now being directed to the development of improved carrier molecules that contain carrier peptides comprising CD4+ T helper cell (Th) epitopes, but which do not possess T-cell suppressive (Ts) functions (Etlinger *et al.* 1990). Peptides which retain only helper functions (CD4+ epitopes) are most suitable as carriers, since their effect is sufficient to induce T cell help but the carrier is small enough to limit or to completely avoid production of anti-carrier antibodies.

Various publications demonstrate the ability of such peptides to confer T-cell help to haptens when covalently linked to them (Etlinger, 1990; Valmori 1992; Sadd 1992; Kumar 1992; Kaliyaperumal, 1995; De Velasco, 1995 and Bixler 1989). However, to date, these publications have not resulted in the development of effective vaccines. There thus remains a great need for the development of new, improved vaccine strategies that are effective in combating diseases caused by encapsulated bacteria in infants and young children.

15

SUMMARY OF THE INVENTION

According to the present invention, there is provided a carrier protein comprising at least five CD4+ T-cell epitopes. Preferably, the carrier protein is conjugated to a polysaccharide. These compounds are useful as immunogenic compounds that may in turn be useful as components of protective vaccines against diseases caused by bacterial pathogens.

A carrier protein is an antigenic polypeptide entity that induces the formation of antibodies directed against an antigen conjugated to it, by the immune system of an organism into which the carrier-antigen conjugate is introduced. The necessity to use carrier proteins results from the fact that although many short epitopes are protective, they are poorly immunogenic. This negates the usefulness of these epitopes in the generation of new and efficacious vaccines. By conjugating an immunogenic carrier protein to a molecule that is non-immunogenic, it is possible to confer the high immunogenicity of the carrier protein onto the conjugate molecule. Such conjugate molecules stimulate the generation of an immune response against the non-immunogenic portion of the conjugate molecule and thus have been effectively used in

vaccines that protect against pathogens for which protective immunity could not otherwise be generated.

Hence, highly immunogenic proteins (such as tetanus toxoid) have historically been used as carriers in order to induce a Th cell response that provides help to B cells for the production of antibodies directed against non-immunogenic epitopes. However, overall effectiveness has not been generally achieved with this approach, since the antibody response to a hapten (the epitope) coupled to a carrier protein can be inhibited when the recipient host has been previously immunised with the unmodified carrier protein. This phenomenon is termed epitope-specific suppression and has now been studied in a variety of hapten-carrier systems.

Coupling of bacterial polysaccharides to carrier proteins has been shown to improve the immunogenicity of the polysaccharide and results in antigens with novel characteristics. Furthermore, the coupling of a thymus-independent (TI) polysaccharide to a protein makes the polysaccharide thymus-dependent (TD).

A CD4⁺ T cell epitope is a peptide epitope that stimulates the activity of those T cells that are MHC Class II restricted. This subset of T cells includes Th cells. Many CD4⁺ epitopes are well known to those of skill in the art and have been shown to confer T cell help to haptens when covalently attached to them (Etlinger *et al*, 1990; Valmori 1992; Sadd 1992; Kumar 1992; Kaliyaperumal, 1995).

The CD4⁺ T epitopes used in the carrier proteins of the present invention ideally comprise peptides that are of as short a length as possible. The epitope will thus retain its characteristics to a sufficient degree to induce T-cell help, yet will be small enough that excessive production of anti-carrier antibodies will be minimised. This is preferable, since it is thought that suppression of immune responses by pre-existing antibodies to carrier epitopes is likely to become a problem in the future if the number of congregate vaccines containing common carrier proteins keeps growing. Furthermore, the use of short peptides as carrier epitopes affords the rational selection of suitable Th epitopes, whilst avoiding stretches of sequence that contain B-cell or T-

suppressor epitopes that will be detrimental to the function of the protein in eliciting a TI immune response.

Suitable proteins from which CD4⁺ epitopes may be selected include tetanus toxin (TT), *Plasmodium falciparum* circumsporozoite, hepatitis B surface antigen, hepatitis B nuclear core protein, *H. influenzae* matrix protein, *H. influenzae* haemagglutinin, diphtheria toxoid, diphtheria toxoid mutant CRM197, group B *N. meningitidis* outer membrane protein complex (OMPC), the pneumococcal toxin pneumolysin, and heat shock proteins from *Mycobacterium bovis*. Other suitable carrier peptide epitopes will be well known to those of skill in the art. The CD4⁺ T-cell epitopes selected from these antigens are recognised by human CD4⁺ T cells.

It has been found that the number of T-cell epitopes present in the carrier protein has a significant influence in conferring T-cell help to oligosaccharide molecules conjugated thereto. The polyepitope carrier protein should contain five or more CD4⁺ T-cell epitopes. Preferably, the polyepitope carrier protein contains between 5 and 20 degenerate CD4⁺ T-cell epitopes, more preferably 5, 6, 7, 8, 9 or 10 degenerate CD4⁺ T-cell epitopes. The use of a number of universal epitopes in the carrier protein has been found to reduce the problem of genetic restriction of the immune response generated against peptide antigens.

In addition to CD4⁺ epitopes, the carrier proteins of the present invention may comprise other peptides or protein fragments, such as epitopes from immunomodulating cytokines such as interleukin-2 (IL-2) or granulocyte-macrophage colony stimulating factor (GM-CSF). Promiscuous peptides (Panina-Bordignon *et al* 1989), the so-called "universal" peptides (Kumar *et al.*, 1992), cluster peptides (Ahlers *et al.*, 1993) or peptides containing both T cell and B cell epitopes (Lett *et al*, 1994) may also be used to recruit various effector systems of the immune system, as required.

The polyepitope carrier protein may be produced by any suitable means, as will be apparent to those of skill in the art. Two preferred methods of construction of carrier proteins according to the invention are direct synthesis and by production of recombinant protein. Preferably, the polyepitope carrier proteins of the present

invention are produced by recombinant means, by expression from an encoding nucleic acid molecule. Recombinant expression has the advantage that the production of the carrier protein is inexpensive, safe, facile and does not involve the use of toxic compounds that may require subsequent removal.

5

When expressed in recombinant form, the carrier proteins of the present invention are generated by expression from an encoding nucleic acid in a host cell. Any host cell may be used, depending upon the individual requirements of a particular vaccine system. Preferably, bacterial hosts are used for the production of recombinant protein, due to the ease with which bacteria may be manipulated and grown. The bacterial host of choice is *Escherichia coli*.

Preferably, if produced recombinantly, the carrier proteins are expressed from plasmids that contain a synthetic nucleic acid insert. Such inserts may be designed by annealing oligonucleotide duplexes that code for the CD4+ T-cell epitopes. The 5' and 3' ends of the synthetic linkers may be designed so as to anneal to each other. This technique allows annealing of the oligonucleotides in a random order, resulting in a mixture of potentially different mini-genes comprising any one of a number of possible combinations of epitopes. This mixture is then cloned into any suitable expression vector and a selection process of expressing clones is then performed. This strategy ensures that only those clones are selected that produce a carrier protein that is not detrimental to the health of the cell in which it is expressed. Conversely, arbitrary selection of the order of epitopes has been found to be less successful.

The ends of the epitope-encoding linkers may be designed so that two codons are introduced between the individual epitopes when annealing takes place. Amino acid residues such as glycine or lysine are examples of suitable residues for use in the spacers. In particular, the use of lysine residues in spacers allows the further congregation of carrier protein to capsular polysaccharide. Additionally, the insertion site in the expression plasmid into which the nucleic acid encoding carrier protein is cloned may allow linkage of the polyepitope carrier protein to a tag, such as the "flag" peptide or polyhistidine. This arrangement facilitates the subsequent purification of recombinant protein.

Nucleic acid encoding the polypeptide carrier protein may be cloned under the control of an inducible promoter, so allowing precise regulation of carrier protein expression. Suitable inducible systems will be well known to those of skill in the art and include the well-known *lac* system (Sambrook *et al.* 1989).

Methods of recombinant expression of carrier proteins according to the invention will be well known to the skilled artisan, but for the purposes of clarity are briefly discussed herein.

10

Mammalian expression systems are known in the art. A mammalian promoter is any DNA sequence capable of binding mammalian RNA polymerase and initiating the downstream (3') transcription of a coding sequence (e.g. structural gene) into mRNA. A promoter will have a transcription initiating region, which is usually placed proximal to the 5' end of the coding sequence, and a TATA box, usually located 25-30 base pairs (bp) upstream of the transcription initiation site. The TATA box is thought to direct RNA polymerase II to begin RNA synthesis at the correct site. A mammalian promoter will also contain an upstream promoter element, usually located within 100 to 200 bp upstream of the TATA box. An upstream promoter element determines the rate at which transcription is initiated and can act in either orientation [Sambrook *et al.* (1989) "Expression of Cloned Genes in Mammalian Cells." In *Molecular Cloning: A Laboratory Manual*, 2nd ed.].

Mammalian viral genes are often highly expressed and have a broad host range; therefore sequences encoding mammalian viral genes provide particularly useful promoter sequences. Examples include the SV40 early promoter, mouse mammary tumour virus LTR promoter, adenovirus major late promoter (Ad MLP), and herpes simplex virus promoter. In addition, sequences derived from non-viral genes, such as the murine metallothionein gene, also provide useful promoter sequences. Expression may be either constitutive or regulated (inducible), depending on the promoter can be induced with glucocorticoid in hormone-responsive cells.

The presence of an enhancer element (enhancer), combined with the promoter elements described above, will usually increase expression levels. An enhancer is a regulatory DNA sequence that can stimulate transcription up to 1000-fold when linked to homologous or heterologous promoters, with synthesis beginning at the normal RNA start site. Enhancers are also active when they are placed upstream or downstream from the transcription initiation site, in either normal or flipped orientation, or at a distance of more than 1000 nucleotides from the promoter [Maniatis *et al.* (1987) *Science* 236:1237; Alberts *et al.* (1989) *Molecular Biology of the Cell*, 2nd ed.]. Enhancer elements derived from viruses may be particularly useful, because they usually have a broader host range. Examples include the SV40 early gene enhancer [Dijkema *et al.* (1985) *EMBO J.* 4:761] and the enhancer/promoters derived from the long terminal repeat (LTR) of the Rous Sarcoma Virus [Gorman *et al.* (1982b) *Proc. Natl. Acad. Sci.* 79:6777] and from human cytomegalovirus [Boshart *et al.* (1985) *Cell* 41:521]. Additionally, some enhancers are regulatable and become active only in the presence of an inducer, such as a hormone or metal ion [Sassone-Corsi and Borelli (1986) *Trends Genet.* 2:215; Maniatis *et al.* (1987) *Science* 236:1237].

A DNA molecule may be expressed intracellularly in mammalian cells. A promoter sequence may be directly linked with the DNA molecule, in which case the first amino acid at the N-terminus of the recombinant protein will always be a methionine, which is encoded by the ATG start codon. If desired, the N-terminus may be cleaved from the protein by *in vitro* incubation with cyanogen bromide.

Alternatively, foreign proteins can also be secreted from the cell into the growth media by creating chimeric DNA molecules that encode a fusion protein comprised of a leader sequence fragment that provides for secretion of the foreign protein in mammalian cells. Preferably, there are processing sites encoded between the leader fragment and the foreign gene that can be cleaved either *in vivo* or *in vitro*. The leader sequence fragment usually encodes a signal peptide comprised of hydrophobic amino acids which direct the secretion of the protein from the cell. The adenovirus tripartite leader is an example of a leader sequence that provides for secretion of a foreign protein in mammalian cells.

Usually, transcription termination and polyadenylation sequences recognised by mammalian cells are regulatory regions located 3' to the translation stop codon and thus, together with the promoter elements, flank the coding sequence. The 3' terminus of the mature mRNA is formed by site-specific post-transcriptional cleavage and polyadenylation [Birnstiel *et al.* (1985) *Cell* 41:349; Proudfoot and Whitelaw (1988) "Termination and 3' end processing of eukaryotic RNA. In *Transcription and splicing* (ed. B.D. Hames and D.M. Glover); Proudfoot (1989) *Trends Biochem. Sci.* 14:105]. These sequences direct the transcription of an mRNA which can be translated into the polypeptide encoded by the DNA. Examples of transcription terminator/polyadenylation signals include those derived from SV40 [Sambrook *et al.* (1989) "Expression of cloned genes in cultured mammalian cells." In *Molecular Cloning: A Laboratory Manual*].

Some genes may be expressed more efficiently when introns (also called intervening sequences) are present. Several cDNAs, however, have been efficiently expressed from vectors that lack splicing signals (also called splice donor and acceptor sites) [see e.g., Gothing and Sambrook (1981) *Nature* 293:620]. Introns are intervening noncoding sequences within a coding sequence that contain splice donor and acceptor sites. They are removed by a process called "splicing," following polyadenylation of the primary transcript [Nevins (1983) *Annu. Rev. Biochem.* 52:441; Green (1986) *Annu. Rev. Genet.* 20:671; Padgett *et al.* (1986) *Annu. Rev. Biochem.* 55:1119; Krainer and Maniatis (1988) "RNA splicing." In *Transcription and splicing* (ed. B.D. Hames and D.M. Glover)].

Usually, the above-described components, comprising a promoter, polyadenylation signal, and transcription termination sequence are put together into expression constructs. Enhancers, introns with functional splice donor and acceptor sites, and leader sequences may also be included in an expression construct, if desired. Expression constructs are often maintained in a replicon, such as an extrachromosomal element (e.g., plasmids) capable of stable maintenance in a host, such as mammalian cells or bacteria. Mammalian replication systems include those derived from animal viruses, which require trans-acting factors to replicate. For example, plasmids containing the replication systems of papovaviruses, such as SV40 [Gluzman (1981) *Cell* 23:175] or polyomavirus, replicate to extremely high copy number in the presence of the

appropriate viral T antigen. Additional examples of mammalian replicons include those derived from bovine papillomavirus and Epstein-Barr virus. Additionally, the replicon may have two replicaton systems, thus allowing it to be maintained, for example, in mammalian cells for expression and in a prokaryotic host for cloning and amplification.

- 5 Examples of such mammalian-bacteria shuttle vectors include pMT2 [Kaufman *et al.* (1989) *Mol. Cell. Biol.* 9:946 and pHEBO [Shimizu *et al.* (1986) *Mol. Cell. Biol.* 6:1074].

10 The transformation procedure used depends upon the host to be transformed. Methods for introduction of heterologous polynucleotides into mammalian cells are known in the art and include dextran-mediated transfection, calcium phosphate precipitation, polybrene mediated transfection, protoplast fusion, electroporation, encapsulation of the polynucleotide(s) in liposomes, and direct microinjection of the DNA into nuclei.

- 15 Mammalian cell lines available as hosts for expression are known in the art and include many immortalised cell lines available from the American Type Culture Collection (ATCC), including but not limited to, Chinese hamster ovary (CHO) cells, HeLa cells, baby hamster kidney (BHK) cells, monkey kidney cells (COS), human hepatocellular carcinoma cells (e.g., Hep G2), and a number of other cell lines.

20

The polynucleotide encoding the protein can also be inserted into a suitable insect expression vector, and is operably linked to the control elements within that vector. Vector construction employs techniques that are known in the art. Generally, the components of the expression system include a transfer vector, usually a bacterial
25 plasmid, which contains both a fragment of the baculovirus genome, and a convenient restriction site for insertion of the heterologous gene or genes to be expressed; a wild type baculovirus with a sequence homologous to the baculovirus-specific fragment in the transfer vector (this allows for the homologous recombination of the heterologous gene in to the baculovirus genome); and appropriate insect host cells and growth media.

30

After inserting the DNA sequence encoding the protein into the transfer vector, the vector and the wild type viral genome are transfected into an insect host cell where the

vector and viral genome are allowed to recombine. The packaged recombinant virus is expressed and recombinant plaques are identified and purified. Materials and methods for baculovirus/insect cell expression systems are commercially available in kit form from, *inter alia*, Invitrogen, San Diego CA ("MaxBac" kit). These techniques are
5 generally known to those skilled in the art and fully described in Summers and Smith, *Texas Agricultural Experiment Station Bulletin No. 1555* (1987) (hereinafter "Summers and Smith").

Prior to inserting the DNA sequence encoding the protein into the baculovirus genome,
10 the above described components, comprising a promoter, leader (if desired), coding sequence of interest, and transcription termination sequence, are usually assembled into an intermediate transplacement construct (transfer vector). This construct may contain a single gene and operably linked regulatory elements; multiple genes, each with its own set of operably linked regulatory elements; or multiple genes, regulated by the
15 same set of regulatory elements. Intermediate transplacement constructs are often maintained in a replicon, such as an extrachromosomal element (e.g., plasmids) capable of stable maintenance in a host, such as a bacterium. The replicon will have a replication system, thus allowing it to be maintained in a suitable host for cloning and amplification.

20

Currently, the most commonly used transfer vector for introducing foreign genes into AcNPV is pAc373. Many other vectors, known to those of skill in the art, have also been designed. These include, for example, pVL985 (which alters the polyhedrin start codon from ATG to ATT, and which introduces a BamHI cloning site 32 basepairs
25 downstream from the ATT; see Luckow and Summers, *Virology* (1989) 17:31.

The plasmid usually also contains the polyhedrin polyadenylation signal (Miller *et al.* (1988) *Ann. Rev. Microbiol.*, 42:177) and a prokaryotic ampicillin-resistance (*amp*) gene and origin of replication for selection and propagation in *E. coli*.

30

Baculovirus transfer vectors usually contain a baculovirus promoter. A baculovirus promoter is any DNA sequence capable of binding a baculovirus RNA polymerase and

initiating the downstream (5' to 3') transcription of a coding sequence (e.g. structural gene) into mRNA. A promoter will have a transcription initiation region which is usually placed proximal to the 5' end of the coding sequence. This transcription initiation region usually includes an RNA polymerase binding site and a transcription initiation site. A baculovirus transfer vector may also have a second domain called an enhancer, which, if present, is usually distal to the structural gene. Expression may be either regulated or constitutive.

Structural genes, abundantly transcribed at late times in a viral infection cycle, provide particularly useful promoter sequences. Examples include sequences derived from the gene encoding the viral polyhedron protein, Friesen *et al.*, (1986) "The Regulation of Baculovirus Gene Expression," in: *The Molecular Biology of Baculoviruses* (ed. Walter Doerfler); EPO Publ. Nos. 127 839 and 155 476; and the gene encoding the p10 protein, Vlak *et al.*, (1988), *J. Gen. Virol.* 69:765.

DNA encoding suitable signal sequences can be derived from genes for secreted insect or baculovirus proteins, such as the baculovirus polyhedrin gene (Carbonell *et al.* (1988) *Gene*, 73:409). Alternatively, since the signals for mammalian cell posttranslational modifications (such as signal peptide cleavage, proteolytic cleavage, and phosphorylation) appear to be recognised by insect cells, and the signals required for secretion and nuclear accumulation also appear to be conserved between the invertebrate cells and vertebrate cells, leaders of non-insect origin, such as those derived from genes encoding human γ -interferon, Maeda *et al.*, (1985), *Nature* 315:592; human gastrin-releasing peptide, Lebacqz-Verheyden *et al.*, (1988), *Molec. Cell. Biol.* 8:3129; human IL-2, Smith *et al.*, (1985) *Proc. Nat'l Acad. Sci. USA*, 82:8404; mouse IL-3, (Miyajima *et al.*, (1987) *Gene* 58:273; and human glucocerebrosidase, Martin *et al.* (1988) *DNA*, 7:99, can also be used to provide for secretion in insects.

A recombinant polypeptide or polyprotein may be expressed intracellularly or, if it is expressed with the proper regulatory sequences, it can be secreted. Good intracellular expression of non-fused foreign proteins usually requires heterologous genes that ideally have a short leader sequence containing suitable translation initiation signals

preceding an ATG start signal. If desired, methionine at the N-terminus may be cleaved from the mature protein by *in vitro* incubation with cyanogen bromide.

Alternatively, recombinant polyproteins or proteins which are not naturally secreted can be secreted from the insect cell by creating chimeric DNA molecules that encode a fusion protein comprised of a leader sequence fragment that provides for secretion of the foreign protein in insects. The leader sequence fragment usually encodes a signal peptide comprised of hydrophobic amino acids which direct the translocation of the protein into the endoplasmic reticulum.

After insertion of the DNA sequence and/or the gene encoding the expression product precursor of the protein, an insect cell host is co-transformed with the heterologous DNA of the transfer vector and the genomic DNA of wild type baculovirus - usually by co-transfection. The promoter and transcription termination sequence of the construct will usually comprise a 2-5kb section of the baculovirus genome. Methods for introducing heterologous DNA into the desired site in the baculovirus virus are known in the art. (See Summers and Smith *supra*; Ju *et al.* (1987); Smith *et al.*, *Mol. Cell. Biol.* (1983) 3:2156; and Luckow and Summers (1989)). For example, the insertion can be into a gene such as the polyhedrin gene, by homologous double crossover recombination; insertion can also be into a restriction enzyme site engineered into the desired baculovirus gene. Miller *et al.*, (1989), *Bioessays* 4:91. The DNA sequence, when cloned in place of the polyhedrin gene in the expression vector, is flanked both 5' and 3' by polyhedrin-specific sequences and is positioned downstream of the polyhedrin promoter.

The newly formed baculovirus expression vector is subsequently packaged into an infectious recombinant baculovirus. Homologous recombination occurs at low frequency (between about 1% and about 5%); thus, the majority of the virus produced after cotransfection is still wild-type virus. Therefore, a method is necessary to identify recombinant viruses. An advantage of the expression system is a visual screen allowing recombinant viruses to be distinguished. The polyhedrin protein, which is produced by the native virus, is produced at very high levels in the nuclei of infected cells at late

times after viral infection. Accumulated polyhedrin protein forms occlusion bodies that also contain embedded particles. These occlusion bodies, up to 15 μ m in size, are highly refractile, giving them a bright shiny appearance that is readily visualised under the light microscope. Cells infected with recombinant viruses lack occlusion bodies. To distinguish recombinant virus from wild-type virus, the transfection supernatant is 5
 10
 15
 20
 25
 30
 35
 40
 45
 50
 55
 60
 65
 70
 75
 80
 85
 90
 95
 100
 105
 110
 115
 120
 125
 130
 135
 140
 145
 150
 155
 160
 165
 170
 175
 180
 185
 190
 195
 200
 205
 210
 215
 220
 225
 230
 235
 240
 245
 250
 255
 260
 265
 270
 275
 280
 285
 290
 295
 300
 305
 310
 315
 320
 325
 330
 335
 340
 345
 350
 355
 360
 365
 370
 375
 380
 385
 390
 395
 400
 405
 410
 415
 420
 425
 430
 435
 440
 445
 450
 455
 460
 465
 470
 475
 480
 485
 490
 495
 500
 505
 510
 515
 520
 525
 530
 535
 540
 545
 550
 555
 560
 565
 570
 575
 580
 585
 590
 595
 600
 605
 610
 615
 620
 625
 630
 635
 640
 645
 650
 655
 660
 665
 670
 675
 680
 685
 690
 695
 700
 705
 710
 715
 720
 725
 730
 735
 740
 745
 750
 755
 760
 765
 770
 775
 780
 785
 790
 795
 800
 805
 810
 815
 820
 825
 830
 835
 840
 845
 850
 855
 860
 865
 870
 875
 880
 885
 890
 895
 900
 905
 910
 915
 920
 925
 930
 935
 940
 945
 950
 955
 960
 965
 970
 975
 980
 985
 990
 995
 1000
 1005
 1010
 1015
 1020
 1025
 1030
 1035
 1040
 1045
 1050
 1055
 1060
 1065
 1070
 1075
 1080
 1085
 1090
 1095
 1100
 1105
 1110
 1115
 1120
 1125
 1130
 1135
 1140
 1145
 1150
 1155
 1160
 1165
 1170
 1175
 1180
 1185
 1190
 1195
 1200
 1205
 1210
 1215
 1220
 1225
 1230
 1235
 1240
 1245
 1250
 1255
 1260
 1265
 1270
 1275
 1280
 1285
 1290
 1295
 1300
 1305
 1310
 1315
 1320
 1325
 1330
 1335
 1340
 1345
 1350
 1355
 1360
 1365
 1370
 1375
 1380
 1385
 1390
 1395
 1400
 1405
 1410
 1415
 1420
 1425
 1430
 1435
 1440
 1445
 1450
 1455
 1460
 1465
 1470
 1475
 1480
 1485
 1490
 1495
 1500
 1505
 1510
 1515
 1520
 1525
 1530
 1535
 1540
 1545
 1550
 1555
 1560
 1565
 1570
 1575
 1580
 1585
 1590
 1595
 1600
 1605
 1610
 1615
 1620
 1625
 1630
 1635
 1640
 1645
 1650
 1655
 1660
 1665
 1670
 1675
 1680
 1685
 1690
 1695
 1700
 1705
 1710
 1715
 1720
 1725
 1730
 1735
 1740
 1745
 1750
 1755
 1760
 1765
 1770
 1775
 1780
 1785
 1790
 1795
 1800
 1805
 1810
 1815
 1820
 1825
 1830
 1835
 1840
 1845
 1850
 1855
 1860
 1865
 1870
 1875
 1880
 1885
 1890
 1895
 1900
 1905
 1910
 1915
 1920
 1925
 1930
 1935
 1940
 1945
 1950
 1955
 1960
 1965
 1970
 1975
 1980
 1985
 1990
 1995
 2000
 2005
 2010
 2015
 2020
 2025
 2030
 2035
 2040
 2045
 2050
 2055
 2060
 2065
 2070
 2075
 2080
 2085
 2090
 2095
 2100
 2105
 2110
 2115
 2120
 2125
 2130
 2135
 2140
 2145
 2150
 2155
 2160
 2165
 2170
 2175
 2180
 2185
 2190
 2195
 2200
 2205
 2210
 2215
 2220
 2225
 2230
 2235
 2240
 2245
 2250
 2255
 2260
 2265
 2270
 2275
 2280
 2285
 2290
 2295
 2300
 2305
 2310
 2315
 2320
 2325
 2330
 2335
 2340
 2345
 2350
 2355
 2360
 2365
 2370
 2375
 2380
 2385
 2390
 2395
 2400
 2405
 2410
 2415
 2420
 2425
 2430
 2435
 2440
 2445
 2450
 2455
 2460
 2465
 2470
 2475
 2480
 2485
 2490
 2495
 2500
 2505
 2510
 2515
 2520
 2525
 2530
 2535
 2540
 2545
 2550
 2555
 2560
 2565
 2570
 2575
 2580
 2585
 2590
 2595
 2600
 2605
 2610
 2615
 2620
 2625
 2630
 2635
 2640
 2645
 2650
 2655
 2660
 2665
 2670
 2675
 2680
 2685
 2690
 2695
 2700
 2705
 2710
 2715
 2720
 2725
 2730
 2735
 2740
 2745
 2750
 2755
 2760
 2765
 2770
 2775
 2780
 2785
 2790
 2795
 2800
 2805
 2810
 2815
 2820
 2825
 2830
 2835
 2840
 2845
 2850
 2855
 2860
 2865
 2870
 2875
 2880
 2885
 2890
 2895
 2900
 2905
 2910
 2915
 2920
 2925
 2930
 2935
 2940
 2945
 2950
 2955
 2960
 2965
 2970
 2975
 2980
 2985
 2990
 2995
 3000
 3005
 3010
 3015
 3020
 3025
 3030
 3035
 3040
 3045
 3050
 3055
 3060
 3065
 3070
 3075
 3080
 3085
 3090
 3095
 3100
 3105
 3110
 3115
 3120
 3125
 3130
 3135
 3140
 3145
 3150
 3155
 3160
 3165
 3170
 3175
 3180
 3185
 3190
 3195
 3200
 3205
 3210
 3215
 3220
 3225
 3230
 3235
 3240
 3245
 3250
 3255
 3260
 3265
 3270
 3275
 3280
 3285
 3290
 3295
 3300
 3305
 3310
 3315
 3320
 3325
 3330
 3335
 3340
 3345
 3350
 3355
 3360
 3365
 3370
 3375
 3380
 3385
 3390
 3395
 3400
 3405
 3410
 3415
 3420
 3425
 3430
 3435
 3440
 3445
 3450
 3455
 3460
 3465
 3470
 3475
 3480
 3485
 3490
 3495
 3500
 3505
 3510
 3515
 3520
 3525
 3530
 3535
 3540
 3545
 3550
 3555
 3560
 3565
 3570
 3575
 3580
 3585
 3590
 3595
 3600
 3605
 3610
 3615
 3620
 3625
 3630
 3635
 3640
 3645
 3650
 3655
 3660
 3665
 3670
 3675
 3680
 3685
 3690
 3695
 3700
 3705
 3710
 3715
 3720
 3725
 3730
 3735
 3740
 3745
 3750
 3755
 3760
 3765
 3770
 3775
 3780
 3785
 3790
 3795
 3800
 3805
 3810
 3815
 3820
 3825
 3830
 3835
 3840
 3845
 3850
 3855
 3860
 3865
 3870
 3875
 3880
 3885
 3890
 3895
 3900
 3905
 3910
 3915
 3920
 3925
 3930
 3935
 3940
 3945
 3950
 3955
 3960
 3965
 3970
 3975
 3980
 3985
 3990
 3995
 4000
 4005
 4010
 4015
 4020
 4025
 4030
 4035
 4040
 4045
 4050
 4055
 4060
 4065
 4070
 4075
 4080
 4085
 4090
 4095
 4100
 4105
 4110
 4115
 4120
 4125
 4130
 4135
 4140
 4145
 4150
 4155
 4160
 4165
 4170
 4175
 4180
 4185
 4190
 4195
 4200
 4205
 4210
 4215
 4220
 4225
 4230
 4235
 4240
 4245
 4250
 4255
 4260
 4265
 4270
 4275
 4280
 4285
 4290
 4295
 4300
 4305
 4310
 4315
 4320
 4325
 4330
 4335
 4340
 4345
 4350
 4355
 4360
 4365
 4370
 4375
 4380
 4385
 4390
 4395
 4400
 4405
 4410
 4415
 4420
 4425
 4430
 4435
 4440
 4445
 4450
 4455
 4460
 4465
 4470
 4475
 4480
 4485
 4490
 4495
 4500
 4505
 4510
 4515
 4520
 4525
 4530
 4535
 4540
 4545
 4550
 4555
 4560
 4565
 4570
 4575
 4580
 4585
 4590
 4595
 4600
 4605
 4610
 4615
 4620
 4625
 4630
 4635
 4640
 4645
 4650
 4655
 4660
 4665
 4670
 4675
 4680
 4685
 4690
 4695
 4700
 4705
 4710
 4715
 4720
 4725
 4730
 4735
 4740
 4745
 4750
 4755
 4760
 4765
 4770
 4775
 4780
 4785
 4790
 4795
 4800
 4805
 4810
 4815
 4820
 4825
 4830
 4835
 4840
 4845
 4850
 4855
 4860
 4865
 4870
 4875
 4880
 4885
 4890
 4895
 4900
 4905
 4910
 4915
 4920
 4925
 4930
 4935
 4940
 4945
 4950
 4955
 4960
 4965
 4970
 4975
 4980
 4985
 4990
 4995
 5000
 5005
 5010
 5015
 5020
 5025
 5030
 5035
 5040
 5045
 5050
 5055
 5060
 5065
 5070
 5075
 5080
 5085
 5090
 5095
 5100
 5105
 5110
 5115
 5120
 5125
 5130
 5135
 5140
 5145
 5150
 5155
 5160
 5165
 5170
 5175
 5180
 5185
 5190
 5195
 5200
 5205
 5210
 5215
 5220
 5225
 5230
 5235
 5240
 5245
 5250
 5255
 5260
 5265
 5270
 5275
 5280
 5285
 5290
 5295
 5300
 5305
 5310
 5315
 5320
 5325
 5330
 5335
 5340
 5345
 5350
 5355
 5360
 5365
 5370
 5375
 5380
 5385
 5390
 5395
 5400
 5405
 5410
 5415
 5420
 5425
 5430
 5435
 5440
 5445
 5450
 5455
 5460
 5465
 5470
 5475
 5480
 5485
 5490
 5495
 5500
 5505
 5510
 5515
 5520
 5525
 5530
 5535
 5540
 5545
 5550
 5555
 5560
 5565
 5570
 5575
 5580
 5585
 5590
 5595
 5600
 5605
 5610
 5615
 5620
 5625
 5630
 5635
 5640
 5645
 5650
 5655
 5660
 5665
 5670
 5675
 5680
 5685
 5690
 5695
 5700
 5705
 5710
 5715
 5720
 5725
 5730
 5735
 5740
 5745
 5750
 5755
 5760
 5765
 5770
 5775
 5780
 5785
 5790
 5795
 5800
 5805
 5810
 5815
 5820
 5825
 5830
 5835
 5840
 5845
 5850
 5855
 5860
 5865
 5870
 5875
 5880
 5885
 5890
 5895
 5900
 5905
 5910
 5915
 5920
 5925
 5930
 5935
 5940
 5945
 5950
 5955
 5960
 5965
 5970
 5975
 5980
 5985
 5990
 5995
 6000
 6005
 6010
 6015
 6020
 6025
 6030
 6035
 6040
 6045
 6050
 6055
 6060
 6065
 6070
 6075
 6080
 6085
 6090
 6095
 6100
 6105
 6110
 6115
 6120
 6125
 6130
 6135
 6140
 6145
 6150
 6155
 6160
 6165
 6170
 6175
 6180
 6185
 6190
 6195
 6200
 6205
 6210
 6215
 6220
 6225
 6230
 6235
 6240
 6245
 6250
 6255
 6260
 6265
 6270
 6275
 6280
 6285
 6290
 6295
 6300
 6305
 6310
 6315
 6320
 6325
 6330
 6335
 6340
 6345
 6350
 6355
 6360
 6365
 6370
 6375
 6380
 6385
 6390
 6395
 6400
 6405
 6410
 6415
 6420
 6425
 6430
 6435
 6440
 6445
 6450
 6455
 6460
 6465
 6470
 6475
 6480
 6485
 6490
 6495
 6500
 6505
 6510
 6515
 6520
 6525
 6530
 6535
 6540
 6545
 6550
 6555
 6560
 6565
 6570
 6575
 6580
 6585
 6590
 6595
 6600
 6605
 6610
 6615
 6620
 6625
 6630
 6635
 6640
 6645
 6650
 6655
 6660
 6665
 6670
 6675
 6680
 6685
 6690
 6695
 6700
 6705
 6710
 6715
 6720
 6725
 6730
 6735
 6740
 6745
 6750
 6755
 6760
 6765
 6770
 6775
 6780
 6785
 6790
 6795
 6800
 6805
 6810
 6815
 6820
 6825
 6830
 6835
 6840
 6845
 6850
 6855
 6860
 6865
 6870
 6875
 6880
 6885
 6890
 6895
 6900
 6905
 6910
 6915
 6

electrophoresis; density gradient centrifugation; solvent extraction, or the like. As appropriate, the product may be further purified, as required, so as to remove substantially any insect proteins which are also secreted in the medium or result from lysis of insect cells, so as to provide a product which is at least substantially free of host debris, e.g., proteins, lipids and polysaccharides.

In order to obtain protein expression, recombinant host cells derived from the transformants are incubated under conditions which allow expression of the recombinant protein encoding sequence. These conditions will vary, dependent upon the host cell selected. However, the conditions are readily ascertainable to those of ordinary skill in the art, based upon what is known in the art.

Bacterial expression techniques are known in the art. A bacterial promoter is any DNA sequence capable of binding bacterial RNA polymerase and initiating the downstream (3") transcription of a coding sequence (e.g. structural gene) into mRNA. A promoter will have a transcription initiation region which is usually placed proximal to the 5' end of the coding sequence. This transcription initiation region usually includes an RNA polymerase binding site and a transcription initiation site. A bacterial promoter may also have a second domain called an operator, that may overlap an adjacent RNA polymerase binding site at which RNA synthesis begins. The operator permits negative regulated (inducible) transcription, as a gene repressor protein may bind the operator and thereby inhibit transcription of a specific gene. Constitutive expression may occur in the absence of negative regulatory elements, such as the operator. In addition, positive regulation may be achieved by a gene activator protein binding sequence, which, if present is usually proximal (5') to the RNA polymerase binding sequence. An example of a gene activator protein is the catabolite activator protein (CAP), which helps initiate transcription of the lac operon in *Escherichia coli* (*E. coli*) [Raibaud *et al.* (1984) *Annu. Rev. Genet.* 18:173]. Regulated expression may therefore be either positive or negative, thereby either enhancing or reducing transcription.

30

Sequences encoding metabolic pathway enzymes provide particularly useful promoter sequences. Examples include promoter sequences derived from sugar metabolising

- enzymes, such as galactose, lactose (*lac*) [Chang *et al.* (1977) *Nature* 198:1056], and maltose. Additional examples include promoter sequences derived from biosynthetic enzymes such as tryptophan (*trp*) [Goeddel *et al.* (1980) *Nuc. Acids Res.* 8:4057; Yelverton *et al.* (1981) *Nucl. Acids Res.* 9:731; U.S. Patent No. 4,738,921; EPO Publ. Nos. 036 776 and 121 775]. The g-laotamase (*bla*) promoter system [Weissmann (1981) "The cloning of interferon and other mistakes." In *Interferon 3* (ed. I. Gresser)], bacteriophage lambda PL [Shimatake *et al.* (1981) *Nature* 292:128] and T5 [U.S. Patent No. 4,689,406] promoter systems also provide useful promoter sequences.
- 10 In addition, synthetic promoters that do not occur in nature also function as bacterial promoters. For example, transcription activation sequences of one bacterial or bacteriophage promoter may be joined with the operon sequences of another bacterial or bacteriophage promoter, creating a synthetic hybrid promoter [U.S. Patent No. 4,551,433]. For example, the *tac* promoter is a hybrid *trp-lac* promoter comprised
- 15 of both *trp* promoter and *lac* operon sequences that is regulated by the *lac* repressor [Amann *et al.* (1983) *Gene* 25:167; de Boer *et al.* (1983) *Proc. Natl. Acad. Sci.* 80:21]. Furthermore, a bacterial promoter can include naturally occurring promoters of non-bacterial origin that have the ability to bind bacterial RNA polymerase and initiate transcription. A naturally occurring promoter of non-bacterial origin can also be
- 20 coupled with a compatible RNA polymerase to produce high levels of expression of some genes in prokaryotes. The bacteriophage T7 RNA polymerase/promoter system is an example of a coupled promoter system [Studier *et al.* (1986) *J. Mol. Biol.* 189:113; Tabor *et al.* (1985) *Proc Natl. Acad. Sci.* 82:1074]. In addition, a hybrid promoter can also be comprised of a bacteriophage promoter and an *E. coli* operator region (EPO
- 25 Publ. No. 267 851).

In addition to a functioning promoter sequence, an efficient ribosome binding site is also useful for the expression of foreign genes in prokaryotes. In *E. coli*, the ribosome binding site is called the Shine-Dalgarno (SD) sequence and includes an initiation

30 codon (ATG) and a sequence 3-9 nucleotides in length located 3-11 nucleotides upstream of the initiation codon [Shine *et al.* (1975) *Nature* 254:34]. The SD sequence is thought to promote binding of mRNA to the ribosome by the pairing of bases between the SD sequence and the 3' end of *E. coli* 16S rRNA [Steitz *et al.* (1979)

"Genetic signals and nucleotide sequences in messenger RNA." In *Biological Regulation and Development: Gene Expression* (ed. R.F. Goldberger)]. To express eukaryotic genes and prokaryotic genes with weak ribosome-binding site [Sambrook *et al.* (1989) "Expression of cloned genes in *Escherichia coli*." In *Molecular Cloning: A Laboratory Manual*].

A DNA molecule may be expressed intracellularly. A promoter sequence may be directly linked with the DNA molecule, in which case the first amino acid at the N-terminus will always be a methionine, which is encoded by the ATG start codon. If desired, methionine at the N-terminus may be cleaved from the protein by *in vitro* incubation with cyanogen bromide or by either *in vivo* or *in vitro* incubation with a bacterial methionine N-terminal peptidase (EPO Publ. No. 219 237).

Fusion proteins provide an alternative to direct expression. Usually, a DNA sequence encoding the N-terminal portion of an endogenous bacterial protein, or other stable protein, is fused to the 5' end of heterologous coding sequences. Upon expression, this construct will provide a fusion of the two amino acid sequences. For example, the bacteriophage lambda cell gene can be linked at the 5' terminus of a foreign gene and expressed in bacteria. The resulting fusion protein preferably retains a site for a processing enzyme (factor Xa) to cleave the bacteriophage protein from the foreign gene [Nagai *et al.* (1984) *Nature* 309:810]. Fusion proteins can also be made with sequences from the *lacZ* [Jia *et al.* (1987) *Gene* 60:197], *trpE* [Allen *et al.* (1987) *J. Biotechnol.* 5:93; Makoff *et al.* (1989) *J. Gen. Microbiol.* 135:11], and *Chey* [EPO Publ. No. 324 647] genes. The DNA sequence at the junction of the two amino acid sequences may or may not encode a cleavable site. Another example is a ubiquitin fusion protein. Such a fusion protein is made with the ubiquitin region that preferably retains a site for a processing enzyme (e.g. ubiquitin specific processing-protease) to cleave the ubiquitin from the foreign protein. Through this method, native foreign protein can be isolated [Miller *et al.* (1989) *Bio/Technology* 7:698].

30

Alternatively, foreign proteins can also be secreted from the cell by creating chimeric DNA molecules that encode a fusion protein comprised of a signal peptide sequence

fragment that provides for secretion of the foreign protein in bacteria [U.S. Patent No. 4,336,336]. The signal sequence fragment usually encodes a signal peptide comprised of hydrophobic amino acids which direct the secretion of the protein from the cell. The protein is either secreted into the growth media (gram-positive bacteria) or into the periplasmic space, located between the inner and outer membrane of the cell (gram-negative bacteria). Preferably there are processing sites, which can be cleaved either *in vivo* or *in vitro* encoded between the signal peptide fragment and the foreign gene.

DNA encoding suitable signal sequences can be derived from genes for secreted bacterial proteins, such as the *E. coli* outer membrane protein gene (*ompA*) [Masui *et al.* (1983), in: *Experimental Manipulation of Gene Expression*; Ghrayeb *et al.* (1984) *EMBO J.* 3:2437] and the *E. coli* alkaline phosphatase signal sequence (*phoA*) [Oka *et al.* (1985) *Proc. Natl. Acad. Sci.* 82:7212]. As an additional example, the signal sequence of the alpha-amylase gene from various *Bacillus* strains can be used to secrete heterologous proteins from *B. subtilis* [Palva *et al.* (1982) *Proc. Natl. Acad. Sci. USA* 79:5582; EPO Publ. No. 244 042].

Usually, transcription termination sequences recognised by bacteria are regulatory regions located 3' to the translation stop codon, and thus together with the promoter flank the coding sequence. These sequences direct the transcription of an mRNA which can be translated into the polypeptide encoded by the DNA. Transcription termination sequences frequently include DNA sequences of about 50 nucleotides capable of forming stem loop structures that aid in terminating transcription. Examples include transcription termination sequences derived from genes with strong promoters, such as the *trp* gene in *E. coli* as well as other biosynthetic genes.

Usually, the above described components, comprising a promoter, signal sequence (if desired), coding sequence of interest, and transcription termination sequence, are put together into expression constructs. Expression constructs are often maintained in a replicon, such as an extrachromosomal element (e.g., plasmids) capable of stable maintenance in a host, such as bacteria. The replicon will have a replication system, thus allowing it to be maintained in a prokaryotic host either for expression or for

cloning and amplification. In addition, a replicon may be either a high or low copy number plasmid. A high copy number plasmid will generally have a copy number ranging from about 5 to about 200, and usually about 10 to about 150. A host containing a high copy number plasmid will preferably contain at least about 10, and more preferably at least about 20 plasmids. Either a high or low copy number vector may be selected, depending upon the effect of the vector and the foreign protein on the host.

Alternatively, the expression constructs can be integrated into the bacterial genome with an integrating vector. Integrating vectors usually contain at least one sequence homologous to the bacterial chromosome that allows the vector to integrate. Integrations appear to result from recombinations between homologous DNA in the vector and the bacterial chromosome. For example, integrating vectors constructed with DNA from various *Bacillus* strains integrate into the *Bacillus* chromosome (EPO Publ. No. 127 328). Integrating vectors may also be comprised of bacteriophage or transposon sequences.

Usually, extrachromosomal and integrating expression constructs may contain selectable markers to allow for the selection of bacterial strains that have been transformed. Selectable markers can be expressed in the bacterial host and may include genes which render bacteria resistant to drugs such as ampicillin, chloramphenicol, erythromycin, kanamycin (neomycin), and tetracycline [Davies *et al.* (1978) *Annu. Rev. Microbiol.* 32:469]. Selectable markers may also include biosynthetic genes, such as those in the histidine, tryptophan, and leucine biosynthetic pathways.

Alternatively, some of the above described components can be put together in transformation vectors. Transformation vectors are usually comprised of a selectable marker that is either maintained in a replicon or developed into an integrating vector, as described above.

Expression and transformation vectors, either extra-chromosomal replicons or integrating vectors, have been developed for transformation into many bacteria. For example, expression vectors have been developed for. *inter alia*, the following bacteria:

Bacillus subtilis [Palva *et al.* (1982) *Proc. Natl. Acad. Sci. USA* 79:5582; EPO Publ. Nos. 036 259 and 063 953; PCT Publ. No. WO 84/04541], Escherichia coli [Shimatake *et al.* (1981) *Nature* 292:128; Amann *et al.* (1985) *Gene* 40:183; Studier *et al.* (1986) *J. Mol. Biol.* 189:113; EPO Publ. Nos. 036 776, 136 829 and 136 907], Streptococcus cremoris [Powell *et al.* (1988) *Appl. Environ. Microbiol.* 54:655]; Streptococcus lividans [Powell *et al.* (1988) *Appl. Environ. Microbiol.* 54:655], Streptomyces lividans [U.S. Patent No. 4,745,056].

Methods of introducing exogenous DNA into bacterial hosts are well-known in the art, and usually include either the transformation of bacteria treated with CaCl₂ or other agents, such as divalent cations and DMSO. DNA can also be introduced into bacterial cells by electroporation. Transformation procedures usually vary with the bacterial species to be transformed. See e.g., [Masson *et al.* (1989) *FEMS Microbiol. Lett.* 60:273; Palva *et al.* (1982) *Proc. Natl. Acad. Sci. USA* 79:5582; EPO Publ. Nos. 036 259 and 063 953; PCT Publ. No. WO 84/04541, Bacillus], [Miller *et al.* (1988) *Proc. Natl. Acad. Sci.* 85:856; Wang *et al.* (1990) *J. Bacteriol.* 172:949, Campylobacter], [Cohen *et al.* (1973) *Proc. Natl. Acad. Sci.* 69:2110; Dower *et al.* (1988) *Nucleic Acids Res.* 16:6127; Kushner (1978) "An improved method for transformation of Escherichia coli with ColE1-derived plasmids. In *Genetic Engineering: Proceedings of the International Symposium on Genetic Engineering* (eds. H.W. Boyer and S. Nicosia); Mandel *et al.* (1970) *J. Mol. Biol.* 53:159; Taketo (1988) *Biochim. Biophys. Acta* 949:318; Escherichia], [Chassy *et al.* (1987) *FEMS Microbiol. Lett.* 44:173 Lactobacillus]; [Fiedler *et al.* (1988) *Anal. Biochem* 170:38, Pseudomonas]; [Augustin *et al.* (1990) *FEMS Microbiol. Lett.* 66:203, Staphylococcus], [Barany *et al.* (1980) *J. Bacteriol.* 144:698; Harlander (1987) "Transformation of Streptococcus lactis by electroporation, in: *Streptococcal Genetics* (ed. J. Ferretti and R. Curtiss III); Perry *et al.* (1981) *Infect. Immun.* 32:1295; Powell *et al.* (1988) *Appl. Environ. Microbiol.* 54:655; Somkuti *et al.* (1987) *Proc. 4th Eur. Cong. Biotechnology* 1:412, Streptococcus].

30

Yeast expression systems are also known to one of ordinary skill in the art. A yeast promoter is any DNA sequence capable of binding yeast RNA polymerase and initiating the downstream (3') transcription of a coding sequence (e.g. structural gene) into

mRNA. A promoter will have a transcription initiation region which is usually placed proximal to the 5' end of the coding sequence. This transcription initiation region usually includes an RNA polymerase binding site (the "TATA Box") and a transcription initiation site. A yeast promoter may also have a second domain called an upstream
 5 activator sequence (UAS), which, if present, is usually distal to the structural gene. The UAS permits regulated (inducible) expression. Constitutive expression occurs in the absence of a UAS. Regulated expression may be either positive or negative, thereby either enhancing or reducing transcription.

- 10 Yeast is a fermenting organism with an active metabolic pathway, therefore sequences encoding enzymes in the metabolic pathway provide particularly useful promoter sequences. Examples include alcohol dehydrogenase (ADH) (EPO Publ. No. 284 044), enolase, glucokinase, glucose-6-phosphate isomerase, glyceraldehyde-3-phosphate-dehydrogenase (GAP or GAPDH), hexokinase, phosphofructokinase, 3-
 15 phosphoglycerate mutase, and pyruvate kinase (PyK) (EPO Publ. No. 329 203). The yeast *PHO5* gene, encoding acid phosphatase, also provides useful promoter sequences [Myanohara *et al.* (1983) *Proc. Natl. Acad. Sci. USA* 80:1].

- In addition, synthetic promoters which do not occur in nature also function as yeast
 20 promoters. For example, UAS sequences of one yeast promoter may be joined with the transcription activation region of another yeast promoter, creating a synthetic hybrid promoter. Examples of such hybrid promoters include the ADH regulatory sequence linked to the GAP transcription activation region (U.S. Patent Nos. 4,876,197 and 4,880,734). Other examples of hybrid promoters include promoters which consist of the
 25 regulatory sequences of either the *ADH2*, *GAL4*, *GAL10*, OR *PHO5* genes, combined with the transcriptional activation region of a glycolytic enzyme gene such as GAP or PyK (EPO Publ. No. 164 556). Furthermore, a yeast promoter can include naturally occurring promoters of non-yeast origin that have the ability to bind yeast RNA
 — polymerase and initiate transcription. Examples of such promoters include, *inter alia*,
 30 [Cohen *et al.* (1980) *Proc. Natl. Acad. Sci. USA* 77:1078; Henikoff *et al.* (1981) *Nature* 283:835; Hollenberg *et al.* (1981) *Curr. Topics Microbiol. Immunol.* 96:119; Hollenberg *et al.* (1979) "The Expression of Bacterial Antibiotic Resistance Genes in the Yeast *Saccharomyces cerevisiae*," in: *Plasmids of Medical, Environmental and*

Commercial Importance (eds. K.N. Timmis and A. Puhler); Mércerau-Puigalon *et al.* (1980) *Gene* 11:163; Panthier *et al.* (1980) *Curr. Genet.* 2:109;].

A DNA molecule may be expressed intracellularly in yeast. A promoter sequence may be directly linked with the DNA molecule, in which case the first amino acid at the N-terminus of the recombinant protein will always be a methionine, which is encoded by the ATG start codon. If desired, methionine at the N-terminus may be cleaved from the protein by *in vitro* incubation with cyanogen bromide.

- 5 Fusion proteins provide an alternative for yeast expression systems, as well as in mammalian, baculovirus, and bacterial expression systems. Usually, a DNA sequence encoding the N-terminal portion of an endogenous yeast protein, or other stable protein, is fused to the 5' end of heterologous coding sequences. Upon expression, this construct will provide a fusion of the two amino acid sequences. For example, the yeast or human superoxide dismutase (SOD) gene, can be linked at the 5' terminus of a foreign gene and expressed in yeast. The DNA sequence at the junction of the two amino acid sequences may or may not encode a cleavable site. See e.g., EPO Publ. No. 196 056. Another example is a ubiquitin fusion protein. Such a fusion protein is made with the ubiquitin region that preferably retains a site for a processing enzyme (e.g. ubiquitin-specific processing protease) to cleave the ubiquitin from the foreign protein. Through this method, therefore, native foreign protein can be isolated (*eg.* WO88/024066).

- Alternatively, foreign proteins can also be secreted from the cell into the growth media by creating chimeric DNA molecules that encode a fusion protein comprised of a leader sequence fragment that provide for secretion in yeast of the foreign protein. Preferably, there are processing sites encoded between the leader fragment and the foreign gene that can be cleaved either *in vivo* or *in vitro*. The leader sequence fragment usually encodes a signal peptide comprised of hydrophobic amino acids which direct the secretion of the protein from the cell.

30

DNA encoding suitable signal sequences can be derived from genes for secreted yeast proteins, such as the yeast invertase gene (EPO Publ. No. 012 873; JPO Publ. No.

62,096,086) and the A-factor gene (U.S. Patent No. 4,588,684). Alternatively, leaders of non-yeast origin, such as an interferon leader, exist that also provide for secretion in yeast (EPO Publ. No. 060 057).

- 5 A preferred class of secretion leaders are those that employ a fragment of the yeast alpha-factor gene, which contains both a "pre" signal sequence, and a "pro" region. The types of alpha-factor fragments that can be employed include the full-length pre-pro alpha factor leader (about 83 amino acid residues) as well as truncated alpha-factor leaders (usually about 25 to about 50 amino acid residues) (U.S. Patent Nos. 4,546,083
10 and 4,870,008; EPO Publ. No. 324 274). Additional leaders employing an alpha-factor leader fragment that provides for secretion include hybrid alpha-factor leaders made with a presequence of a first yeast, but a pro-region from a second yeast alphafactor. (See e.g., PCT Publ. No. WO 89/02463.)
- 15 Usually, transcription termination sequences recognised by yeast are regulatory regions located 3' to the translation stop codon, and thus together with the promoter flank the coding sequence. These sequences direct the transcription of an mRNA which can be translated into the polypeptide encoded by the DNA. Examples of transcription terminator sequence and other yeast-recognised termination sequences, such as those
20 coding for glycolytic enzymes.

- Usually, the above described components, comprising a promoter, leader (if desired), coding sequence of interest, and transcription termination sequence, are put together into expression constructs. Expression constructs are often maintained in a replicon,
25 such as an extrachromosomal element (e.g., plasmids) capable of stable maintenance in a host, such as yeast or bacteria. The replicon may have two replication systems, thus allowing it to be maintained, for example, in yeast for expression and in a prokaryotic host for cloning and amplification. Examples of such yeast-bacteria shuttle vectors include YEp24 [Botstein *et al.* (1979) *Gene* 8:17-24], pCl/1 [Brake *et al.* (1984) *Proc.*
30 *Natl. Acad. Sci USA* 81:4642-4646], and YRp17 [Stinchcomb *et al.* (1982) *J. Mol. Biol.* 158:157]. In addition, a replicon may be either a high or low copy number plasmid. A high copy number plasmid will generally have a copy number ranging from about 5 to

about 200, and usually about 10 to about 150. A host containing a high copy number plasmid will preferably have at least about 10, and more preferably at least about 20. Enter a high or low copy number vector may be selected, depending upon the effect of the vector and the foreign protein on the host. See e.g., Brake *et al.*, *supra*.

5

Alternatively, the expression constructs can be integrated into the yeast genome with an integrating vector. Integrating vectors usually contain at least one sequence homologous to a yeast chromosome that allows the vector to integrate, and preferably contain two homologous sequences flanking the expression construct. Integrations appear to result from recombinations between homologous DNA in the vector and the yeast chromosome [Orr-Weaver *et al.* (1983) *Methods in Enzymol.* 101:228-245]. An integrating vector may be directed to a specific locus in yeast by selecting the appropriate homologous sequence for inclusion in the vector. See Orr-Weaver *et al.*, *supra*. One or more expression construct may integrate, possibly affecting levels of recombinant protein produced [Rine *et al.* (1983) *Proc. Natl. Acad. Sci. USA* 80:6750]. The chromosomal sequences included in the vector can occur either as a single segment in the vector, which results in the integration of the entire vector, or two segments homologous to adjacent segments in the chromosome and flanking the expression construct in the vector, which can result in the stable integration of only the expression construct.

20

Usually, extrachromosomal and integrating expression constructs may contain selectable markers to allow for the selection of yeast strains that have been transformed. Selectable markers may include biosynthetic genes that can be expressed in the yeast host, such as *ADE2*, *HIS4*, *LEU2*, *TRP1*, and *ALG7*, and the G418 resistance gene, which confer resistance in yeast cells to tunicamycin and G418, respectively. In addition, a suitable selectable marker may also provide yeast with the ability to grow in the presence of toxic compounds, such as metal. For example, the presence of *CUP1* allows yeast to grow in the presence of copper ions [Butt *et al.* (1987) *Microbiol. Rev.* 51:351].

30

Alternatively, some of the above described components can be put together into transformation vectors. Transformation vectors are usually comprised of a selectable marker that is either maintained in a replicon or developed into an integrating vector, as described above.

5

Expression and transformation vectors, either extrachromosomal replicons or integrating vectors, have been developed for transformation into many yeasts. For example, expression vectors have been developed for, *inter alia*, the following yeasts: *Candida albicans* [Kurtz, *et al.* (1986) *Mol. Cell. Biol.* 6:142], *Candida maltosa* [Kunze, *et al.* (1985) *J. Basic Microbiol.* 25:141], *Hansenula polymorpha* [Gleeson, *et al.* (1986) *J. Gen. Microbiol.* 132:3459; Roggenkamp *et al.* (1986) *Mol. Gen. Genet.* 202:302], *Kluyveromyces fragilis* [Das, *et al.* (1984) *J. Bacteriol.* 158:1165], *Kluyveromyces lactis* [De Louvencourt *et al.* (1983) *J. Bacteriol.* 154:737; Van den Berg *et al.* (1990) *Bio/Technology* 8:135], *Pichia guillermondii* [Kunze *et al.* (1985) *J. Basic Microbiol.* 25:141], *Pichia pastoris* [Cregg, *et al.* (1985) *Mol. Cell. Biol.* 5:3376; U.S. Patent Nos. 4,837,148 and 4,929,555], *Saccharomyces cerevisiae* [Hinnen *et al.* (1978) *Proc. Natl. Acad. Sci. USA* 75:1929; Ito *et al.* (1983) *J. Bacteriol.* 153:163], *Schizosaccharomyces pombe* [Beach and Nurse (1981) *Nature* 300:706], and *Yarrowia lipolytica* [Davidow, *et al.* (1985) *Curr. Genet.* 10:380471 Gaillardin, *et al.* (1985) *Curr. Genet.* 10:49].

Methods of introducing exogenous DNA into yeast hosts are well-known in the art, and usually include either the transformation of spheroplasts or of intact yeast cells treated with alkali cations. Transformation procedures usually vary with the yeast species to be transformed. See e.g., [Kurtz *et al.* (1986) *Mol. Cell. Biol.* 6:142; Kunze *et al.* (1985) *J. Basic Microbiol.* 25:141; *Candida*]; [Gleeson *et al.* (1986) *J. Gen. Microbiol.* 132:3459; Roggenkamp *et al.* (1986) *Mol. Gen. Genet.* 202:302; *Hansenula*]; [Das *et al.* (1984) *J. Bacteriol.* 158:1165; De Louvencourt *et al.* (1983) *J. Bacteriol.* 154:1165; Van den Berg *et al.* (1990) *Bio/Technology* 8:135; *Kluyveromyces*]; [Cregg *et al.* (1985) *Mol. Cell. Biol.* 5:3376; Kunze *et al.* (1985) *J. Basic Microbiol.* 25:141; U.S. Patent Nos. 4,837,148 and 4,929,555; *Pichia*]; [Hinnen *et al.* (1978) *Proc. Natl. Acad. Sci. USA* 75:1929; Ito *et al.* (1983) *J. Bacteriol.* 153:163 *Saccharomyces*]; [Beach and Nurse

(1981) *Nature* 300:706; Schizosaccharomyces]; [Davidow *et al.* (1985) *Curr. Genet.* 10:39; Gaillardin *et al.* (1985) *Curr. Genet.* 10:49; Yarrowia].

Methods for the isolation and purification of recombinant proteins will be well known to those of skill in the art and are summarised, for example in Sambrook *et al.* (1989). Particularly in bacteria such as *E. coli*, the recombinant protein will form inclusion bodies within the bacterial cell, thus facilitating its preparation. If produced in inclusion bodies, the carrier protein may need to be refolded to its natural conformation. Methods for renaturing proteins to their natural folded state are well known in the art.

Species in which the carrier proteins of the present invention may be immunogenic and thus effective in eliciting an immune response include all mammals, especially humans. In most cases, it will be preferred that the carrier proteins of the present invention are active eliciting an immune response in humans. The population of humans that are in greatest need of protection from disease caused by encapsulated bacteria are infants of between approximately 3 months and 2 years of age. It is during this period that the infants generally do not receive protection from mothers' milk and do not yet possess a sufficiently well-developed immune system themselves to generate an immune response against polysaccharide antigens.

According to a further aspect of the present invention, there are also provided nucleic acid molecules encoding carrier proteins according to the first aspect of the invention. As will be apparent to the skilled artisan, such nucleic acid molecules will be designed using the genetic code so as to encode the epitope that is desired.

Additionally, in order to precisely tailor the exact properties of the encoded carrier proteins, the skilled artisan will appreciate that changes may be made at the nucleotide level from known epitope sequences, by addition, substitution, deletion or insertion of one or more nucleotides. Site-directed mutagenesis (SDM) is the method of preference used to generate mutated carrier proteins according to the present invention. There are many techniques of SDM now known to the skilled artisan, including oligonucleotide-directed mutagenesis using PCR as set out, for example by Sambrook *et al.*, (1989) or using commercially available kits.

Most carrier proteins produced by such techniques of mutagenesis will be less efficacious than wild type proteins. However, it may be that in a minority of cases, such changes produce molecules with improved carrier protein function as desired, such as
5 greater immunogenicity in a certain organism.

The nucleic acid molecules according to this aspect of the present invention may comprise DNA, RNA or cDNA and may additionally comprise nucleotide analogues in the coding sequence. Preferably, the nucleic acid molecules will comprise DNA.

10

A further aspect of the present invention provides a host cell containing a nucleic acid encoding a carrier protein. A still further aspect provides a method comprising introducing the encoding nucleic acid into a host cell or organism. Introduction of nucleic acid may employ any available technique. In eukaryotic cells, suitable
15 techniques may include calcium phosphate transfection, DNA-dextran, electroporation, liposome-mediated transfection or transduction using retrovirus or other viruses such as vaccinia. In bacterial cells, suitable techniques may include calcium chloride transformation, electroporation or transfection using bacteriophage. Introduction of the nucleic acid may be followed by causing or allowing expression from the nucleic acid,
20 for example by culturing host cells under conditions for allowing expression of the gene.

In one embodiment, the nucleic acid is integrated into the genome of the host cell. Integration may be promoted by the inclusion of sequences that promote recombination
25 with the genome, in accordance with standard techniques (see Sambrook *et al.*, 1989).

According to a further embodiment of the present invention, there is provided a carrier protein comprising at least five CD4+ T-cell epitopes, conjugated to polysaccharide. By polysaccharide is meant any linear or branched polymer consisting of
30 monosaccharide residues, usually linked by glycosidic linkages, and thus includes oligosaccharides. Preferably, the polysaccharide will contain between 2 and 50 monosaccharide units, more preferably between 6 and 30 monosaccharide units.

The polysaccharide component may be based on or derived from polysaccharide components of the polysaccharide capsule from many Gram positive and Gram negative bacterial pathogens such as *H. influenzae*, *N. meningitidis* and *S. pneumoniae*. This capsule represents a major virulence factor that is important for nasopharyngeal colonisation and systemic invasion. Other bacteria from which polysaccharide components may be conjugated to the carrier proteins of the present invention include *Staphylococcus aureus*, *Klebsiella*, *Pseudomonas*, *Salmonella typhi*, *Pseudomonas aeruginosa*, and *Shigella dysenteriae*. Polysaccharide components suitable for use according to this aspect of the present invention include the Hib oligosaccharide, lipopolysaccharide from *Pseudomonas aeruginosa* (Seid and Sadoff, 1981), lipopolysaccharides from *Salmonella* (Konadu *et al.*, 1996) and the O-specific polysaccharide from *Shigella dysenteriae* (Chu *et al.*, 1991). Other polysaccharide components suitable for use in accordance with the present invention will be well-known to those of skill in the art.

Fragments of bacterial capsular polysaccharide may be produced by any suitable method, such as by acid hydrolysis or ultrasonic irradiation (Szn *et al.*, 1986). Other methods of preparation of the polysaccharide components will be well known to those of skill in the art.

The polysaccharide component of the conjugate vaccine should preferably be coupled to the carrier protein by a covalent linkage. A particularly preferred method of coupling polysaccharide and protein is by reductive amination. Other methods include: activation of the polysaccharide with cyanogen bromide followed by reaction with adipic acid dihydrazide (spacer) and by conjugation to carboxide groups of carrier protein using soluble carbodiimides (Shneerson *et al.*, 1986); functionalisation of the carrier protein with adipic acid dihydrazide followed by coupling to cyanogen bromide activated polysaccharides (Dick *et al.*, 1989); chemical modification of both the carrier protein and the polysaccharide followed by their coupling (Marburg *et al.*, 1986; Marburg *et al.*, 1987 and 1989). In some cases, polysaccharides containing carboxide groups such as group C meningococcal polysaccharides can be directly conjugated to proteins using soluble carbodiimides. Polysaccharides can also be activated using alternative agents such as CDAP (1-cyano-4-dimethylamino-pyrridinium tetrafluoroborate) and then

directly conjugated to the carrier protein (Konadu *et al*, 1996). Periodate-treated polysaccharides or oligosaccharides can all be conjugated to proteins by means of reductive amination (Jennings and Lugowsky, 1982; Anderson, 1983; Insel, 1986). Alternatively, oligosaccharides obtained by acidic hydrolysis can be chemically
5 derivatised by introducing into their reducing end groups an hydrocarbon spacer bearing an active ester terminus; this activated oligosaccharide can be conjugated to the selected carrier protein (Costantino *et al*, 1992).

The polysaccharide molecule may be coupled to the carrier protein by a spacer
10 molecule, such as adipic acid. This spacer molecule can be used to facilitate the coupling of protein to polysaccharide. After the coupling reaction has been performed, the conjugate may be purified by diafiltration or other known methods to remove unreacted protein or polysaccharide components.

15 According to a further aspect of the present invention there is provided a method of production of a carrier protein according to the first aspect of the present invention, comprising the steps of:

- (a) constructing oligonucleotide molecules that encode peptide epitopes;
- (b) annealing the oligonucleotide molecules to form duplexes;
- 20 (c) introducing the oligonucleotide duplexes into an expression vector so as to encode a fusion protein;
- (d) introducing the expression vector into a bacterial host cell to allow expression of the fusion protein;
- (e) isolating the fusion protein produced from a culture of said bacteria.

25

Optionally, this method may additionally comprise conjugating the carrier protein to a polysaccharide molecule.

Preferably, the bacterial host cell used in this method is an *E. coli* bacterial host cell.

30

According to the further aspect of the present invention, there is provided a composition comprising a carrier protein that contains at least five CD4⁺ T-Cell epitopes conjugated to a polysaccharide, in conjunction with a pharmaceutically acceptable excipient. Such

a composition may be rationally designed so as to provide protection against disease caused by pathogenic bacteria such as *H. influenzae*, *S. pneumoniae*, *N. meningitidis*, *Staphylococcus aureus*, *Klebsiella*, *Pseudomonas* and *S. typhi* and accordingly, may be used as a vaccine. Vaccines according to the invention may either be prophylactic (*ie.* to prevent infection) or therapeutic (*ie.* to treat disease after infection).

By pharmaceutically-acceptable excipient is meant any compound that does not itself induce the production of antibodies harmful to the individual receiving the composition. The excipient should be suitable for oral, subcutaneous, intramuscular, topical or intravenous administration. Suitable compounds are typically large, slowly metabolised macromolecules such as proteins, polysaccharides, polylactic acids, polyglycolic acids, polymeric amino acids, amino acid copolymers, lipid aggregates (such as oil droplets or liposomes) and inactive virus particles. Such compounds are well known to those of skill in the art. Additionally, these compounds may function as immunostimulating agents ("adjuvants"). Furthermore, the antigen may be conjugated to a bacterial toxoid.

Preferred adjuvants to enhance effectiveness of the composition include, but are not limited to: (1) aluminium salts (alum), such as aluminium hydroxide, aluminium phosphate, aluminium sulphate, *etc.*; (2) oil-in-water emulsion formulations (with or without other specific immunostimulating agents such as muramyl peptides or bacterial cell wall components), such as for example (a) MF59™ (WO 90/14837), containing 5% Squalene, 0.5% Tween™ 80, and 0.5% Span 85 (optionally containing various amounts of MTP-PE, although not required) formulated into submicron particles using a microfluidizer (b) SAF, containing 10% Squalene, 0.4% Tween 80, 5% pluronic-blocked polymer L121, and thr-MDP either microfluidised into a submicron emulsion or vortexed to generate a larger particle size emulsion, and (c) Ribi™ adjuvant system (RAS), containing 2% Squalene, 0.2% Tween 80, and one or more bacterial cell wall components from the group consisting of monophosphorylipid A (MPL), trehalose dimycolate (TDM), and cell wall skeleton (CWS), preferably MPL + CWS (Detox™); (3) saponin adjuvants, such as Stimulon™ may be used or particles generated therefrom such as ISCOMs (immunostimulating complexes); (4) Freund's complete and incomplete adjuvants (CFA & IFA); (5) cytokines, such as interleukins (*eg.* IL-1, IL-2, IL-4, IL-5, IL-6, IL-7, IL-12, *etc.*), interferons (*eg.* IFNγ), macrophage colony

stimulating factor (M-CSF), tumor necrosis factor (TNF), etc; and (6) other substances that act as immunostimulating agents to enhance the efficacy of the composition. Alum and MF59™ are preferred.

- 5 As mentioned above, muramyl peptides include, but are not limited to, N-acetyl-muramyl-L-threonyl-D-isoglutamine (thr-MDP), N-acetyl-normuramyl-L-alanyl-D-isoglutamine (nor-MDP), N-acetylmuramyl-L-alanyl-D-isoglutaminyl-L-alanine-2-(1'-2'-dipalmitoyl-*sn*-glycero-3-hydroxyphosphoryloxy)-ethylamine (MTP-PE), *etc.*
- 10 The immunogenic compositions (*eg.* the antigen, pharmaceutically acceptable carrier, and adjuvant) typically will contain diluents, such as water, saline, glycerol, ethanol, *etc.* Additionally, auxiliary substances, such as wetting or emulsifying agents, pH buffering substances, and the like, may be present in such vehicles.
- 15 Typically, the immunogenic compositions are prepared as injectables, either as liquid solutions or suspensions; solid forms suitable for solution in, or suspension in, liquid vehicles prior to injection may also be prepared. The preparation also may be emulsified or encapsulated in liposomes for enhanced adjuvanticity effect, as discussed above.
- 20 Immunogenic compositions used as vaccines comprise an immunologically effective amount of the carrier protein, as well as any other of the above-mentioned components, as needed. By "immunologically effective amount", it is meant that the administration of that amount to an individual, either in a single dose or as part of a series, is effective for treatment or prevention. This amount varies depending upon the health and physical
- 25 condition of the individual to be treated, the taxonomic group of individual to be treated (*eg.* non-human primate, primate, *etc.*), the capacity of the individual's immune system to synthesise antibodies, the degree of protection desired, the formulation of the vaccine, the treating doctor's assessment of the medical situation, and other relevant factors. It is expected that the amount will fall in a relatively broad range that can be
- 30 determined through routine trials.

The immunogenic compositions are conventionally administered parenterally *eg.* by injection. either subcutaneously or intramuscularly. They may also be administered to

mucosal surfaces (eg. oral or intranasal), or in the form of pulmonary formulations, suppositories, or transdermal applications. Dosage treatment may be a single dose schedule or a multiple dose schedule. The vaccine may be administered in conjunction with other immunoregulatory agents.

5

As an alternative to protein-based vaccines, DNA vaccination may be employed [eg. Robinson & Torres (1997) *Seminars in Immunology* 9:271-283; Donnelly *et al.* (1997) *Annu Rev Immunol* 15:617-648]. Accordingly, rather than comprise a peptide, oligopeptide, or polypeptide compound, the vaccines of the invention might comprise nucleic acid encoding these compounds.

10

According to a further aspect of the invention, there is provided a process for the formulation of an immunogenic composition comprising bringing a carrier protein according to the first aspect of the invention, conjugated to a polysaccharide, into association with a pharmaceutically-acceptable carrier, optionally with an adjuvant.

15

According to a still further aspect of the present invention, there is provided a method of vaccinating a mammal, preferably a human against a disease, comprising administering to the mammal a composition of carrier protein conjugated to polysaccharide, optionally with a pharmaceutically-acceptable carrier such as an adjuvant.

20

Various aspects and embodiments of the present invention will now be described in more detail by way of example, with particular reference to the carrier proteins N6 and N10 conjugated to HIB capsular polysaccharide. It will be appreciated that modification of detail may be made without departing from the scope of the invention. All publications, patents, and patent applications cited herein are incorporated in full by reference.

25

BRIEF DESCRIPTION OF THE FIGURES

30

Figure 1 a schematic representation of the construction of the N6 protein.

Figure 2 illustrates the N6 and N10 constructs and their respective DNA and amino acid sequences. The histidine tag, the flag peptide, the Fxa cutting site and the CD4+ T cell epitopes are underlined.

- 5 Figure 3 is a Coomassie-stained SDS-PAGE gel of total protein extracts prepared from induced *E. coli* clones producing the different polyepitope proteins. Lane A: negative control (TG1 cells containing pQE30 vector with no insert); lane B: TG1 cells containing the pQE30-N10 plasmid; lane C: TOP10 cells containing the pTrc-N10 plasmid; lane D: TOP10 cells containing the pTrc-N6 plasmid; lane E: low molecular
10 weight markers.

Figure 4 is an immunoblot of the SDS-PAGE gel that is illustrated in Figure 3. The Western blot was incubated with a rabbit antiserum specific for the flag peptide and then with a peroxidated anti-rabbit IgG antibody. The immune reaction was then
15 revealed using 4-chloro-1-naphthol as substrate for the peroxidase.

Figure 5 is an SDS-PAGE Coomassie-stained gel containing different samples obtained during the procedure of purification of the N6 protein. Lane A: starting material (total protein of the induced TOP10 *E. coli* cells containing pTrc-N6 plasmid; lane B: soluble
20 proteins (supernatant obtained after centrifugation of the total protein sample); lane C: proteins soluble in 1M urea (supernatant obtained after washing the insoluble proteins with 1M urea); lane D: inclusion bodies (pellet obtained after washing the insoluble proteins with 1M urea); lane E: N6 protein obtained from purification on Ni²⁺ NTA resin using the immobilised metal affinity chromatography (IMAC) technique; lane F:
25 low molecular weight markers.

Figure 6 is an immunoblot of the SDS-PAGE gel that is illustrated in Figure 5. The Western blot was incubated with a rabbit antiserum specific for the flag peptide and then with a peroxidated anti-rabbit IgG antibody. The immune reaction was then
30 revealed using 4-chloro-1-naphthol as substrate for the peroxidase.

Figure 7: Proliferative response of two human T cell clones specific for P30TT (GG-22 clone) and P2TT (KSIMK-140 clone) after stimulation with the respective synthetic

peptides (controls) and with conjugated or nunconjugated polyepitope proteins (cpm: counts per minute).

Figure 8: Peripheral blood mononuclear cells (PBMC) proliferation assay. PBMC from three healthy donors, RR, EB and MC, immune to tetanus toxoid were stimulated with tetanus toxoid, P2TT, N6, N6-Hib and N10-Hib.

10 **DETAILED DESCRIPTION OF THE INVENTION**

MATERIALS AND METHODS

15 **Summary of standard procedures and techniques**

The practice of the present invention will employ, unless otherwise indicated, conventional techniques of molecular biology, microbiology, recombinant DNA, and immunology, which are within the skill of the art. Such techniques are explained fully in the literature *eg.* Sambrook *Molecular Cloning: A Laboratory Manual, Second Edition* (1989); *DNA Cloning, Volumes I and ii* (D.N Glover ed. 1985); *Oligonucleotide Synthesis* (M.J. Gait ed, 1984); *Nucleic Acid Hybridization* (B.D. Hames & S.J. Higgins eds. 1984); *Transcription and Translation* (B.D. Hames & S.J. Higgins eds. 1984); *Animal Cell Culture* (R.I. Freshney ed. 1986); *Immobilised Cells and Enzymes* (IRL Press, 1986); B. Perbal, *A Practical Guide to Molecular Cloning* (1984); the *Methods in Enzymology* series (Academic Press, Inc.), especially volumes 154 & 155; *Gene Transfer Vectors for Mammalian Cells* (J.H. Miller and M.P. Calos eds. 1987, Cold Spring Harbor Laboratory); Mayer and Walker, eds. (1987), *Immunochemical Methods in Cell and Molecular Biology* (Academic Press, London); Scopes, (1987) *Protein Purification: Principles and Practice*, Second Edition (Springer-Verlag, N.Y.), and *Handbook of Experimental Immunology, Volumes I-IV* (D.M. Weir and C. C. Blackwell eds 1986).

Plasmids, strains and T cell clones.

PEMBLex2 plasmid was derived from pEMBL8M (Dente L. and Cortese R. *Meth. Enzymol.* (1987), 155: 111-9) by inserting a λP_L promoter and a polylinker into the
 5 *EcoRI* and *HindIII* sites. The commercial vectors pTrc-His and pQE30 were purchased from Invitrogen and Qiagen respectively. *E. coli* strains used as recipients of the above plasmids were: K12 Δ H1 Δ Trp for pEMBLex2, TOP10 for pTrc-His and TG1 for pQE30.

10 Human T cell clones KSMIK 140 and GG-22 specific for P2TT and P30TT respectively were kindly provided by Dr. A. Lanzavecchia (Basel, Switzerland).

Construction of recombinant plasmids that express the N6 polyepitope carrier protein.

15 Complementary oligodeoxyribonucleotide pairs coding for P2TT, P21TT, P23TT, P30TT1, P32TT and Pft3 T cell epitopes (Table 1) and for a Flag peptide were synthesised using the DNA synthesiser ABI394 (Perkin Elmer) and the reagents from Cruachem (Glasgow, Scotland). The oligo pairs were separately annealed in T4 DNA
 20 ligase buffer (Boehringer Mannheim) and equimolar amounts of each annealing reaction were mixed and ligated at room temperature for 3 hours using T4 DNA ligase (Boehringer Mannheim).

The ligase reaction was then loaded onto a 1% agarose gel and subjected to
 25 electrophoresis. The bands corresponding to the DNA fragments of expected size were isolated, purified and cloned into the pEMBLex2 expression vector using standard protocols (Sambrook *et al.*, 1989). After transformation, a rabbit antiserum specific for the Flag peptide was used to perform colony-screenings (Sambrook *et al.*, 1989) in order to identify recombinant protein producing clones. Protein extracts from positive
 30 clones were analysed using SDS-PAGE to further select for clones on the basis of recombinant protein size.

TABLE 1

P23TT	(1084-1099) VSIDKFRIFCKANPK (Demotz 1993)
P32TT	(1174-1189) LKFIHKRYTPNNEIDS (Demotz 1993)
P21TT	(1064-1079) IREDNNITLKLDRCNN (Dr A. Lanzavecchia, pers. comm.)
PF T3	(380-398) EKKIAKMEKASSVFNVVN (Hammer 1993, Sinigaglia 1988)
P30TT	(947-967) FNNFTVSFWLRVPKVSASHLE (Demotz 1993)
P2TT	(830-843) QYIKANSKFIGITE (Demotz 1993)
HA	(307-319) PKYVKQNTLKLAT (Hammer 1993, O'Sullivan 1991, Alexander 1994)
HBVnc	(50-69) PHHTALRQAILCWGELMTLA (Ferrari 1991, Penna 1992, Alexander 1994)
HBsAG	(19-33) FFLLTRILTIQSLD (Greenstein 1992, Celis 1988)
MT	(17-31) YSGPLKAEIAQRLEDV (Alexander 1994)
Flag peptide	MDYKDDDD

After nucleotide sequencing of the selected clones, a clone named pEMBLN6 was shown to contain six different T cell epitopes with no repetitive sequences. The N6 insert was then PCR-amplified and transferred to pTrc-His expression vector (Invitrogen) using standard techniques (Sambrook *et al.*, 1989). The generation of the N6 expressing plasmids is summarised in Figure 1.

Construction of recombinant plasmids that express the N10 polyepitope carrier protein

10

Using synthetic oligodeoxyribonucleotides and standard cloning techniques (Sambrook *et al.*, 1989), four additional CD4+ T cell epitopes were added to the N6 protein: HBVnc, HA, HbsAg, and MT (Table I). HBVnc and HA were sequentially introduced into pTrc-N6 by means of two consecutive cloning steps; to the resulting plasmid the HbsAg and MT epitopes were added in a single cloning step.

After DNA sequencing, a correct construct (pTrc-N10) coding for the expected ten epitope polyepitope protein was identified. The N10 coding insert was then transferred from pTrc-N10 to pQE30 (Qiagen) by means of PCR. The sequence of the resulting pQE-N10 construct was then confirmed by DNA sequencing.

Purification of the polyepitope carrier proteins.

A TOP10 *E. coli* clone carrying pTrc-N6 plasmid, a TOP10 clone carrying pTrc-N10 plasmid and a TG1 *E. coli* clone carrying pQE-N10 plasmid were grown in 500 ml LB medium containing 100 µg/ml ampicillin at 37°C. At 0.3 OD₆₀₀, the expression of the polyepitope proteins was induced for 3.5 hours by adding 1 mM IPTG. Cells were disrupted by sonication, the insoluble fraction was collected by centrifugation, dissolved with buffer A (6 M guanidinium-HCL, 100 mM NaH₂PO₄, 10 mM Tris base, pH 8) and adsorbed with 2 ml of Ni²⁺NTA resin (Qiagen).

Then, the resin was packed in a column and washed with buffer A. Guanidinium-HCL was removed from the column by washing with buffer B (8 M Urea, 100 mM NaH₂PO₄,

10 mM Tris base) pH 8. After a wash with buffer B pH 6.5, recombinant proteins were eluted with a 20 ml buffer B gradient from pH 6.5 to pH 4. The fractions containing the recombinant polypeptide proteins were pooled together and dialysed against PBS, 10% glycerol, pH 7.2. Proteins were analysed by means of polyacrylamide gels and protein content was assayed by the Bradford method.

Preparation and activation of Hib oligosaccharides.

The Hib capsular polysaccharide can be prepared according to the protocol described in Gotschlich *et al.* (1981) *J. Biol. Chem.* **256**: 8915-8921.

1.99L of a 10 mg/ml solution of Hib polysaccharide was hydrolysed in 0.01M acetic acid at 76°C for 5 hours. After chilling, neutralization and 0.2µm filtration, the resulting oligosaccharide population had an average degree of polymerisation (avDP) of 8 as measured by the chemical ratio between ribose and reducing groups.

NaCl was then added to the hydrolysate until a concentration of 0.16 M was attained, then diluted 1:1 with 0.16M NaCl/ 10mM acetate pH 6 and submitted to tangential flow ultrafiltration on a 10 kDa membrane in order to remove high molecular weight species.

Ultrafiltration comprised approximately 11-fold concentration followed by 15 cycles of diafiltration against 0.16 M NaCl/ 10mM acetate, pH 6. The retentate was discarded. The permeate was diluted :1 with water and 0.22 µm filtered. Chemical analysis revealed an avDp of 8.1.

The permeate obtained from 10 kDa UF was loaded, at a linear flow rate of 150 cm/h, onto a Q-Sepharose Fast Flow column [10 cm (ID); 5,5 cm (h)] equilibrated with 0.08 M NaCl/0.05 M sodium acetate pH 6. After adsorption, low molecular weight fragments (up to 5 repeats) were removed by washing the column with 10 column volumes of equilibrating buffer and then eluted with 3 column volumes of 0.5 M NaCl/0.005M sodium acetate pH 6. The eluate was 0.2 µm filtered and then analysed for avDp and ion exchange analytical chromatography. AvDP resulted at 17.3, ion

exchange analytical chromatography on Mono Q HR 5/5 revealed the absence of any small fragments until DP 5.

To introduce a terminal amino group, reductive amination was then performed; to the
5 fractionated Hib oligosaccharide obtained from Q-Sepharose chromatography, ammonium chloride 35mg/ml and sodium cyanoborohydride 12 mg/ml final concentrations were added. After stirring, the solution was 0.2 μ m filtered and incubated at 37°C for 120 hours. The amino oligosaccharide was then purified from excess of reagents by precipitation with 95° EtOH (81° final concentration) in the cold for 15-20
10 hours. The precipitated oligosaccharide was then recovered by centrifugation, solubilized in NaCl 0.4M using approximately 1/4 of the starting volume and precipitated again at 81° EtOH in the cold for 15-20 hours.

The amino-oligosaccharide was again recovered by centrifugation and solubilized in
15 about 300 ml of 0.02 M NaCl. After having taken a sample for analysis, the resulting solution was then dried using a rotary evaporator.

Colorimetric amino group analysis confirmed the introduction of a primary amino group into the oligosaccharide.

20 Derivatisation to active ester was then performed as follows. The amino-oligosaccharide was solubilised in distilled water at a concentration of 40 μ mol of amino groups per ml. The solution was then diluted 10-fold with DMSO. Triethylamine was added in molar ratio to the amino groups of 2:1. N-hydroxysuccinimido diester of adipic acid was then
25 added in a molar ratio to the amino groups of 12:1. The reaction mixture was kept under gentle stirring for 2 hours at RT. The activated oligosaccharide was then purified from the excess of reagents by precipitation into 10 volumes of 1-4 dioxane under stirring. After 30 minutes in the cold the precipitate was collected onto a sintered glass filter, washed onto the filter with dioxane and then dried under vacuum. The dried
30 activated oligosaccharide was analysed for its content of active ester groups by a colorimetric method; this test showed the presence of 62.1 μ mol of active ester per mg of dried oligosaccharide.

The above-obtained activated oligosaccharide was then used for conjugation experiments.

Conjugation of the polyepitope carrier protein with Hib capsular oligosaccharides and purification of the conjugates.

33.4nmoles of recombinant carrier protein and 669nmoles of activated Hib oligosaccharide in a final volume of 0.5 ml 10mM phosphate buffer, pH 7, were gently stirred overnight at RT and brought up to 5ml 1M $(\text{NH}_4)_2\text{SO}_4$, 10mM phosphate pH7.

10 The sample was subjected to FPLC on a 1ml Phenyl Sepharose 5/5 HR column (Pharmacia). 1ml fractions were collected both during washing (1M $(\text{NH}_4)_2\text{SO}_4$, 10mM phosphate, pH 7) and elution (10mM phosphate, pH 7). Two peaks corresponding to the non-adsorbed material and to the eluted material were obtained. The pooled fractions corresponding to the non-adsorbed material and the pooled fractions

15 corresponding to the elution peak were subjected to protein and ribose content determination and to SDS-PAGE and Western blot analysis.

A protocol to conjugate recombinant proteins to oligosaccharides directly on Ni^{2+} -NTA resin was also developed. Recombinant proteins were purified as described above, but

20 the final dialysis step was omitted. The protein content of the 8M urea fraction pool was measured with the Bradford assay. The pH of the eluted proteins was adjusted to pH8 and adsorption on 1ml pre-equilibrated Ni^{2+} -NTA resin was again performed in a batch mode. Urea was removed by washing with 4x25ml 100mM phosphate buffer pH 7.5. The resin was suspended in 1ml 100mM phosphate buffer pH 7.5 and a 20-fold

25 molar excess of activated Hib oligosaccharide (as compared to the protein that was adsorbed on the resin) was added to the suspension. The mixture was gently stirred overnight at RT packed in a column, and washed with 50ml 100mM phosphate buffer pH 7.5 to remove unconjugated oligosaccharide.

30 Elution of the conjugate was performed with 100mM phosphate buffer pH 4. Peak fractions were pooled and dialysed against PBS, pH 7.2. The conjugate was analysed by Coomassie staining of SDS-PAGE gels and Western immunoblot using an anti-flag

rabbit antibody. The protein/carbohydrate ratio of the glycoconjugate was determined upon Bradford assay and ribose content determination.

Cultures of PBMCs and T cell clones.

5

Culture medium for PBMCs was RPMI 1640 (Gibco Laboratories, Paisley, Scotland) supplemented with 2mM L-glutamine, 1% nonessential amino acids, 1mM sodium pyruvate, gentamycin (50µg/ml), and 5% human serum (RPMI-HS) or 10% foetal calf serum (RPMI-FCS). For the growth of T-cell lines and clones, RPMI-HS was
10 supplemented with 50U of recombinant interleukin-2 (rIL-2: Hoffmann La Roche, Nutley, NJ) per ml.

PBMC Proliferation Assay.

15 Frozen PBMC (10^5) from healthy adults immune to tetanus toxoid were thawed and cultured in duplicate wells of 96-well flat-bottomed microplates, in 0.2ml of RPMI-HS (Di Tommaso *et al*, 1997). The recombinant proteins and tetanus toxoids (Chiron, Siena) were added to wells at the final concentration of 10µg/ml. After 5 days of culture. 1µCi of [3 H] thymidine (specific activity: 5Ci/mmol, Amersham) was added to
20 each well and DNA-incorporated radioactivity was measured after an additional 16hrs by liquid scintillation counting.

Proliferation assay of T cell clones.

25 Two Human T cell clones, KSMIK 140, and GG-22, specific for P2TT and P30TT respectively, and the respective peptides were kindly provided by Dr. A. Lanzavecchia (Basel, Switzerland). T cells (2×10^4) were cultured with autologous irradiated Epstein Barr Virus-transformed B lymphocytes (3×10^4) in 0.2 ml of RPMI-FCS in 96-well flat-bottomed microplates in duplicate wells. Synthetic peptides and conjugated or
30 unconjugated recombinant proteins were added to cultures at a final concentration of 10µg/ml. After 2 days, 1 µCi of [3 H]thymidine was added and the radioactivity incorporated was measured by liquid scintillation counting after an additional 16 hours.

In some experiments, carrier proteins and their conjugates were pre-incubated with APCs for 2-4 hours, then APCs were washed and cultured with T cell clones. This procedure was used to limit possible proteolytic degradation by serum proteases and to be more confident that epitope presentation would be due to intracellularly-processed epitopes.

Immunogenicity tests.

In a first experiment, equal doses of the glycoconjugates and of the polysaccharide (2.5µg as polysaccharide) in presence of 0.5mg of aluminium hydroxide as adjuvant were injected subcutaneously into groups of eight BALB/c and C57BL/6 mice (female, 7-week-old) on days 0, 21 and 35. Mice were bled on day -1 (pre-immune), 20 (pre-2), 34 (pre-3) and 45 (post-3) and individual sera collected and stored at -80°C before ELISA assay.

In a second experiment, equal doses of the glycoconjugates and of the polysaccharide (2.5µg as polysaccharide) in the presence of 0.5mg of aluminium hydroxide as adjuvant were injected subcutaneously into groups of eight Swiss ('D1 and BALB/c mice (female, 7-week-old) on days 0, 10 and 20. A boost of 2.5µg of purified Hib polysaccharide (HibCPS) in presence of 0.5 mg of aluminium hydroxide was then given to each mouse at day 70. Mice were bled on day -1 (pre-immune). 35 (post-vaccination), 68 (pre-boost) and 85 (post-boost) and individual sera collected and stored at -80°C before ELISA assay.

ELISA.

Nunc Maxisorp 96-well flat-bottomed plates were coated by overnight incubation at 4°C with 1µg/ml (as polysaccharide) of a human serum albumin (HSA) and *H. influenzae* type b polysaccharide conjugate (HSA-Hib). After washing, wells were over-coated using 1% (w/v) gelatin in PBS, pH 7.2 for 3 additional hours at 37°C. Serum samples were diluted 1:50 in 5mM phosphate buffer, pH 7.2 containing 75mM NaCL 1% (w/v) BSA and 0.05% (w/v) Tween-20 and dispensed in duplicate into the wells. Sera from untreated mice were pooled and diluted 1:50 as above and dispensed into 8 wells. After

overnight incubation at 4°C, plates were washed three times with 5mM phosphate buffer, pH 7.2 containing 75mM NaCl and 0.05% (w/v) Tween-20. Then, alkaline phosphate-conjugated goat IgG anti-mouse IgG diluted 1:1000 and 5mM phosphate buffer, pH 7.2 containing 75mM NaCl, 1% (w/v) BSA and 0.05% (w/v) Tween-20 were added to each well, and incubated 3 hours at 37°C.

After repeated washing, 100µl of a chromogen-substrate, *p*-nitrophenylphosphate, in a diethylenamine solution, were added to each well. Reaction was stopped after 20min by adding a 4N NaOH solution. Then, the plate was read at 405nm with a reference wavelength of 595nm. Titres were expressed as absorbencies at 405nm (A_{405nm}). Mice were considered responders when the average A_{405nm} was found equal to or higher than four times the average of absorbencies of the eight wells with the sera from untreated animals. According to the European Pharmacopoeia [PA/PH/Exp15/T(93)3ANP] four out of eight mice should be responders.

In the second experiment, mice were considered responders when the average A_{405nm} was found four times the average of the absorbencies of eight pre-immune sera of the same group of treatment.

The anti-carrier response was assayed as above described for anti-Hib response using plates coated with N10 or N6 (coating concentration = 2µg/ml).

RESULTS

Construction of the Polyepitope Carrier Proteins.

Using the approaches described in materials and methods, three clones were selected. One clone produced N6 protein, whilst both the other two clones produced N10 protein. The clone expressing N6 protein comprised the plasmid pTrc-N5+transformed in the Top10 *E. coli* strain. As deduced from nucleotide sequencing, this plasmid coded for a protein having an hexahistidine amino terminal tail followed in sequence by a flag peptide, a *Fxa* site, and the following T cell epitopes: P23TT, P32TT, P21TT, PfCs,

P30TT and P2TT. All the epitopes were spaced by a KG amino acid sequence (Figure 2).

The two clones that produced N10 protein were firstly the Top10 *E.coli* strain containing the plasmid pTrc-N10, and the TG1 *E. Coli* strain containing the plasmid pQE-N10. Both these clones contained the N6 coding sequence fused to a carboxy terminal sequence coding for four additional T cell epitopes which were in the order: HBVnc, HA, HbsAg, and MT (Figure 2).

Protein Expression and Purification.

Figures 3 and 4 depict protein expression of the three synthetic proteins. The addition of four new epitopes (HBVnc, HA, HbsAg, and MT) to N6 in pTrc-His (lane D) to obtain N10 protein (lane C) resulted in a remarkable reduction of protein expression. An attempt to increase the expression level of N10 simply involved changing the expression vector (from pTrc)-His to pQE30) and the *E. coli* strain (from Top10 to TG1). As seen in Figures 3 and 4, the amount of N10 expressed by pQE30-N10 in TG1 (lane B) was notably higher than the same protein expressed by pTrc-N10 (lane C). This is thought possibly to be due to the fact that whereas N6 protein was effectively assembled by the *E. coli* strain in the order of epitopes most suited to the organism, whereas the addition of four further epitopes was effectively forced and thus was less natural. However, the fact that the level of N10 expression was notably increased by simply changing expression vector (from pTrc-His to PQE30) and *E. coli* strain (from TOP-10 to TG1) suggests that additional factors, other than epitope combination, play a role in protein expression.

All clones expressing synthetic protein produced the recombinant proteins mainly in the form of inclusion bodies. Purification of N6 and N10 proteins from inclusion bodies solubilised with 8M urea using an immobilised metal affinity chromatography (IMAC) procedure in the presence of 8M urea resulted in the loss of a high percentage of protein which was elutable with a 6.5-4 pH gradient (data not shown).

On the contrary, almost all of the histidine-tagged protein was eluted with the 6.5-4 pH gradient when starting inclusion bodies were solubilised with 6M guanidine hydrochloride (Figures 5 and 6). Using this protocol 7.8mg of N6 was purified from a litre of culture. The N10 protein that was employed in immunisation and T cell proliferation experiments was purified from pTrc-N10 clone.

Given the lower expression of recombinant protein shown by this clone we decided to purify N10 protein by solubilising whole cells with guanidinium in such a way as to exploit soluble and insoluble (inclusion bodies) proteins for IMAC purification. With this procedure 1.5 mg of purified N10 protein was obtained from a litre of culture. The higher success of solubilisation using 6M guanidinium is thought to be due to the ability of this compound to solubilise the carrier proteins in monomeric form.

Hib oligosaccharide conjugation to polypeptide proteins.

Using the phenyl sepharose FPLC protocol we obtained a purified N6-Hib conjugate having a protein content of 79.4 µg/ml, and an oligosaccharide content of 42.7µg/ml.

We observed that 30% of conjugated protein was unable to bind to phenyl sepharose in the presence of 1M (NH₄)₂SO₄. In addition, 30-40% of carrier protein was previously lost during a dialysis step to remove urea before the conjugation reaction. To overcome these problems it was checked if it was possible to perform the conjugation reactions when the protein was adsorbed on the Ni²⁺-NTA resin. We observed that the Hib oligosaccharide was unable to bind Ni²⁺-NTA resin at any pH, suggesting the feasibility of this approach and predicting that no interference due to the oligosaccharide could influence the elution of the protein once conjugation had taken place.

A reaction was thus set up involving protein adsorption on Ni²⁺-NTA resin in the presence of 8M urea, urea removal, conjugation with oligosaccharide, washing, and conjugate elution. No aggregation phenomena were observed for the eluted conjugate. Using this procedure we obtained a purified N6-Hib conjugate having a protein content of 320µg/ml and an oligosaccharide content of 370µg/ml. and a purified N10-Hib having a protein content of 113µg/ml and an oligosaccharide content of 114µg/ml.

By using a 1:10 protein to carbohydrate molar ratio to conjugate oligosaccharide to recombinant carriers, we observed that a fraction of protein remained unconjugated (as judged by Coomassie staining of SDS-PAGE gel and Western immunoblot; data not shown). When a 1:20 protein to carbohydrate stoichiometric ratio was used, all the purified recombinant proteins were found to be completely conjugated, in fact, by analysing Coomassie-stained gels and western immunoblots using an anti-Flag antibody. We observed that after conjugation of N6 and N10 with Hib oligosaccharides these molecules increased their molecular weight, appearing as a high molecular weight smear, and proteins were no longer visible at the expected molecular weight for N6 and N10 monomers. This suggested that the synthetic proteins were completely conjugated to Hib oligosaccharides (data not shown).

Recognition of carrier proteins and their conjugates by human T lymphocytes.

To investigate whether T cell epitopes contained in the polypeptides were recognised by human T cells we used T cell clones specific for the TT universal epitopes p2TT and p30TT (Demotz *et al.* 1993). Figure 7 shows that N6 is recognised by both clones not only as a simple polypeptide but also after it has been conjugated with polysaccharide. Remarkably, N6-Hib is recognised even better than unconjugated N6 by the T cell clone specific for P2TT. N10-Hib is recognised by the clone specific for p2TT but is poorly recognised by the clone specific for P30TT. In both cases N10-Hib exerts the same stimulatory activity as the synthetic peptide. The N10 clone was not tested in these experiments.

Once assessed that the T cell epitopes contained in the carrier proteins are correctly presented to T lymphocytes, we asked whether these carriers maintain their stimulatory capacity when presented to a heterogeneous population of lymphocytes such as PBMC. This could be predictive of whether our carriers might function as such once injected into subjects immune to antigens whose epitopes are included in the carriers themselves. For this purpose we used PBMC from donors immune to TT (A. Di Tommaso *et al.* 1997), since TT epitopes are the most represented in our polypeptides. Figure 8 shows that all the formulations were able to stimulate PBMC proliferation.

However, the incubation of PBMC with a synthetic peptide representing one of the epitopes included in both N6 and N10 constructs failed to exert a stimulatory effect. As a positive control, the PBMC were also incubated with 10 µg/ml of TT, that in all cases induced a proliferative response. Interestingly, the N6 polyepitope protein turned out to be the most potent PBMC stimulator among those tested in two out of three volunteers.

Immunogenicity Tests.

The carrier effect of the proteins N10 and N6 in comparison with CRM197 was assayed in mice in several glycoconjugate vaccines. Once coupled to Hib oligosaccharides the carrier proteins were injected in different mouse strains to verify the potential of their carrier effect. In BALB/c mice, an equivalent anti-Hib response was found when CRM197 and N10 were used as carrier proteins, whilst a lower response was found when N6 was used as carrier protein. This result was evident when the results were expressed using titres, while responder percentages failed to evidence the lower anti-Hib response obtained with the N6 protein carrier.

In C57BL/6 mice, the N6 protein gave a negative result, while positive results were obtained with CRM197 and N10, even if to a lower extent. These results were evident both using titres or responder percentages to express the results. When the results were expressed as a responder percentage, the high carrier effect of CRM197 and N10 was well evidenced with respect to N6, whose results were lower than 50% at day -34 and day -45 bleedings, after a comparable primary response (pre-2 bleeding, day 20).

Table II reports the results of the experiments in BALB/c and C57BL/6 mice.

TABLE II

BALB/c MICE									
		RESPONDER (%)			A405 x 1000 (GMT's)				
DAY	BLEEDING	N10-Hib	N5+146-Hib	CRM-Hib	N10-Hib	N5+146-Hib	CRM-Hib		
0	PRE-IMMUNE	0	0	0	10	17	12		
20	PRE-2	33.3	33.3	50	135	162	257		
34	POST-2/PRE-3	100	100	100	2022	1356	1969		
45	POST-3	100	100	100	1717	1368	1616		
C57BL/6 MICE									
		RESPONDER (%)			A405 x 1000 (GMT's)				
DAY	BLEEDING	N10-Hib	N5+146-Hib	CRM-Hib	N10-Hib	N5+146-Hib	CRM-Hib		
0	PRE-IMMUNE	0	0	0	28	38	31		
20	PRE-2	83.3	83.3	83.3	136	192	609		
34	POST-2/PRE-3	83.3	33.3	100	1451	306	2612		
45	POST-3	100	33.3	100	1731	222	2240		

In Swiss CDI mice, the titres obtained with the N10 carrier protein were equivalent to those obtained with CRM197. The anti-Hib titres increased after immunisation up to the 70th day, when a polysaccharide boost was given to assay whether or not an immunological memory was induced in the treated mice. No boost effect was observed
5 with any carrier, although when CRM197 or N10 were used as carrier protein the titre did not decrease. In this mouse strain the immunisation with N6-Hib glycoconjugate give results very similar to the controls (polysaccharide and alum). The boost effect was not evidenced even in BALB/c mice that evoke a lower response with respect to Swiss CD1 mice.

10

The results are summarised in Table III.

Immunisation of different mice strains with Hib oligosaccharides conjugated to the artificial carrier proteins resulted in a good carrier effect exerted by N10 ,whilst N6
15 gave unsatisfactory results. This suggests that the size of the protein or the number of T cell epitopes has a high influence in providing T cell help to the oligosaccharides.

TABLE III

SWISS CD1 MICE							
		TITRE GMT's (A _{405nm} x 10 ³)					
DAY	BLEEDING	CRM-Hib	N5+146-Hib	N10-Hib	PsHib	ALUM	
-1	PRE-IMMUNISATION	59	98	156	166	175	
35	POST-IMMUNISATION	1577	471	1007	227	243	
68	PRE-BOOST	2082	889	1789	590	461	
85	POST-BOOST	2073	630	1767	364	479	
SWISS CD1 MICE							
		RESPONDER (%)					
DAY	BLEEDING	CRM-Hib	N5+146-Hib	N10-Hib	PsHib	ALUM	
-1	PRE-IMMUNISATION	0	0	0	0	0	
35	POST-IMMUNISATION	100	50	62.5	0	0	
68	PRE-BOOST	87.5	87.5	100	25	25	
85	POST-BOOST	87.5	62.5	85.7	12.5	37.5	

REFERENCES

- Agadjanyan M, Luo P, Westerink MAJ, Carey LA, Jutchins W, Steplewski Z, Weiner DB, Kieber-Emmons T (1997) Peptide mimicry of carbohydrate epitopes on human immunodeficiency virus. *Nature Biotech.* **15**: 547-551.
- Ahlers J.D., (1993) *J. Immunol.* **150**: 5647-5665.
- Anderson P, Picchichero ME, Insel RA (1985). Immunogens consisting of oligosaccharides from the capsule of *H. influenzae* type b coupled to diphtheria toxoid or the toxin protein CRM₁₉₇ *J Clin Invest* **76**: 52-59.
- Anderson P, Picchichero ME, Insel RA (1985). Immunization of 2-month-old infants with protein-coupled oligosaccharides derived from the capsule of *H. influenzae* type b. *J Pediatr* **107**: 346-351.
- Anderson P. (1983) Antibody responses to *H. Influenzae* type b and diphtheria toxin induced by conjugates of oligosaccharides of the type b capsule with the non-toxic protein CRM₁₉₇. *Infect Immun.* **39**: 233-238
- Andreoni J, Kaythy H, Densen P (1993) Vaccination and the role of capsular polysaccharide antibody in prevention of recurrent meningococcal disease in late complement component-deficient individuals. *J. Infect. Dis.* **168**: 227-231.
- Bixler, G.S. *et al*, (1989) *Adv. Exp. Med. Biol* **V**:175-180.
- Constantino P, Viti S, Podda A, Velmonte M.A., Nencioni L, Rappuoli R (1992). Development and phase 1 clinical testing of a conjugate vaccine against meningococcus A and C. *Vaccine* **10**: 691-698.
- De Velasco EA, Merkus D, Anderton S, Verheul AFM, Lizzio EF, Van der Zee R, Van Eden W, Hoffman T, Vehoef J, Snippe H (1995) Synthetic peptides representing T-cell

epitopes act as carriers in pneumococcal polysaccharide conjugate vaccines. *Infect Immun* 63: 961-968.

- 5 Dick WE, Beurret Mjr. Glycoconjugates of bacterial carbohydrate antigens. A survey and consideration of design and preparation factors, *Conjugate Vaccines* (J.M. Cruse and R.E. Lewis, eds.) Karger, Basel, 1989, p.48.

Etlinger HM, Gillessen D, Lahm HW, Matile H, Schonfeld HJ, Trzeciak A (1990) Use of prior vaccination for the development of new vaccines. *Science* 249: 423-425.

10

Goldblatt D, Levinsky RJ, Turner MW (1992) Role of cell wall polysaccharide in the assessment of IgG antibodies to the capsular polysaccharides of *Streptococcus pneumoniae* in childhood. *J. Infect. Dis.* 166: 632-634.

- 15 Jennings H.J. and C. Lugowsky, Immunogenic conjugates, U.S. Patent No. 4,902,506 (1990).

Holmes SJ, Granoff DM (1992) The biology of *Haemophilus influenzae* type b vaccination failure. *J. Infect. Dis.* 165: S121-S128

20

Insel RA, Anderson PW (1986). Oligosaccharide-protein conjugate vaccines induce and prime for oligoclonal IgG antibody responses to *H. influenzae* b capsular polysaccharide in human infants. *J Exp Med* 163: 262-269

- 25 Jennings HJ, Lugowsky C (1981). Immunochemistry of group A, B, and C meningococcal polysaccharide-tetanus toxoid conjugates. *J. Immunol* 127: 1011-1018.

Kaliyaperumal A, Chauhan VS, Talwar GP Raghupathy R (1995) Carrier-induced epitope-specific regulation at its bypass in a protein-protein conjugate. *Eur J Immunol* 25: 3375-3380.

30

Konadu E, Schiloach G, Bryla D.A., Robins JB, Szu SC (1996) Synthesis, characterization, and immunological properties in mice of conjugates composed of

detoxified lipopolysaccharides of *Salmonella paratyphi* A bound to tetanus toxoid with emphasis on the role of O acetyls. *Infect Immun* **64**: 2709-2715.

- 5 Kumar A, Arora R, Kaur P, Chauhan VS, Sharma P (1992) "Universal" T helper cell determinants enhance immunogenicity of a *Plasmodium falciparum* merozoite surface antigen peptide. *J Immunol* **148**: 1499-1505.

Lett, E. *et al*, (1994) *Infect Immun* 785-792.

- 10 Liptak GS, McConnochie KM, Roghmann KJ, Panzer JA (1997) Decline of pediatric admissions with *Haemophilus influenzae* type b in New York state, 1982 through 1993: Relation to immunisations. *J. Pediatr* **130**: 923-930.

- 15 Marburg S, Jorn D, Tolman RL, Arison B, McCauley J, Kniskern PJ, Hagopian A, Vella PP (1986). Bimolecular chemistry of macromolecules – synthesis of bacterial polysaccharide conjugates with *Neisseria meningitidis* membrane protein. *J. Am Chem Soc* **108**, 5282.

- 20 McNamara MK, Ward RE, Kohler H (1984) Monoclonal idiotope vaccine against *Streptococcus pneumoniae* infection. *Science* **226**: 1325-1326.

Moxon ER and Kroll JS (1990) The role of bacterial polysaccharide capsules as virulence factors. *Curr. Top. Microbiol. Immunol.* **150**: 65-85.

- 25 Panina-Bordignon P, *et al*, (1989) *Eur J Immunol.* **19**: 2237-2242.

- Robbins JB, Schneerson R, Anderson P, Smith DH (1996) Prevention of systemic infections, especially meningitis, caused by *Haemophilus influenzae* type b: impact on public health and implications for other polysaccharide-based vaccines. *JAMA* **276**:
30 1181-1185.

S. Marburg, R.L. Tolman, and P.J. Kniskern, Covalently-modified polyanionic bacterial polysaccharides and immunogenic protein with bigeneric spacers, and methods of

preparing such polysaccharides and conjugates and of confirming covalency, U.S. Patent Nos. 4,695,624 (1987) and 4,882,317 (1989).

5 Sad S, Rao K, Arora R, Talwar GP, Raghupathy R (1992) Bypass of carrier-induced epitope-specific suppression using a T-helper epitope. *Immunology* 76: 599-603.

Schneerson R, Robbins JB, Parke JC, Bell C, Schlesselman JJ, Stton A, Wang Z, Schiffman G, Karpas A, Shiloach J (1986). Quantitative and qualitative analysis of serum antibodies elicited in adults by *Haemophilus influenzae* type b and
10 pneumococcus type 6A capsular polysaccharide-tetanus toxoid conjugates. *Infect Immun* 52: 519.

Tunkel AR and Scheld WM (1993) Pathogenesis and pathophysiology of bacterial meningitis. *Clin. Microbiol. Rev.* 6: 118-136.

15

Valmori D, Pessi A, Bianchi E, Corradin G (1992) Use of human universally antigenic tetanus toxin T cell epitopes as carriers for human vaccination. *J Immunol* 149: 717-721.

20

CLAIMS

- 1 A carrier protein comprising at least five CD4+ T cell epitopes.
- 5 2 A carrier protein according to claim 1, wherein the CD4+ epitopes are derived from a pathogenic bacterium or virus.
- 3 A carrier protein according to claim 1 or 2, wherein the CD4+ epitopes are derived from tetanus toxin, *Plasmodium falciparum* circumsporozoite protein, hepatitis B
10 surface antigen, hepatitis B nuclear core protein, influenza matrix protein, influenza haemagglutinin, diphtheria toxoid, diphtheria toxin mutant CRM 197, group B *Neisseria meningitidis* outer membrane protein complex or pertussis toxin.
- 4 A carrier protein according to any one of the preceding claims wherein the
15 CD4+ epitopes are selected from the P23TT, P32TT, P21TT, PfCs, P30TT, P2TT, HBVnc, HA, HbsAg and MT CD4+ epitopes.
- 5 A carrier protein according to claim 1, that comprises the P23TT, P32TT, P21TT, PfCs, P30TT, P2TT, HBVnc, HA, HbsAg and MT CD4+ epitopes.
20
- 6 A carrier protein according to claim 1, that comprises the P23TT, P32TT, P21TT, PfCs, P30TT and P2TT CD4+ epitopes.
- 7 A carrier protein according any one of the preceding claims, wherein the CD4+
25 epitopes are human CD4+ epitopes.
- 8 A carrier protein according to any one of the preceding claims in an oligomeric form.
- 30 9 A carrier protein according to any one of the preceding claims, conjugated to polysaccharide.

10 A carrier protein according to claim 9, wherein the polysaccharide is a *Haemophilus influenzae* type B polysaccharide.

11 A carrier protein according to claim 9, wherein the polysaccharide is derived
5 from any one of the following organisms: *S. pneumoniae*, *N. meningitidis*, *S. aureus*, *Klebsiella*, or *S. typhimurium*.

12 A carrier protein according to any one of claims 9-11 where the polysaccharide is conjugated to protein by a covalent linkage.

10

13 A carrier protein according to claim any one of claims 9-11, wherein the polysaccharide is conjugated to protein by reductive amination.

14 A carrier protein according to any one of claims 9-13, wherein there are between
15 two and ten protein units for each polysaccharide unit.

15 A carrier protein according to any one of claims 1 to 14 for use as a pharmaceutical.

20 16 Use of the carrier protein according to any one of claims 1 to 14 as a pharmaceutical.

17 The carrier protein according to any one of claims 1 to 14 for use as a vaccine or as a component of a vaccine.

25

18 Use of a carrier protein according to any one of claims 1 to 14 as a vaccine or vaccine component.

19 A vaccine comprising a carrier protein according to any one of claims 1 to 14.

30

20 A method of production of vaccination comprising introducing into a mammal, preferably a human, a carrier protein according to any one of claims 1 to 14.

- 21 The carrier protein according to any one of claims 1 to 14 for use as a protective immunogen in the control of diseases caused by encapsulated bacteria.
- 22 A nucleic acid molecule which encodes a carrier protein according to any one of
5 claims 1 to 8.
- 23 The nucleic acid molecule of claim 22 which comprises DNA.
- 24 A cloning or expression vector comprising a nucleic acid molecule according to
10 either of claims 22-23.
- 25 A host cell transformed or transfected with the vector of claim 24.
- 26 A transgenic animal that has been transformed by a nucleic acid molecule
15 according to either of claims 22 or 23 or by a vector according to claim 24.
- 27 A method of preparing a carrier protein according to any one of claims 1 to 6,
comprising expressing a vector according to claim 24 in a host cell and culturing said host
cell under conditions where said protein is expressed, and recovering said protein thus
20 expressed.
- 28 A method of production of a carrier protein according to any one of claims 1-6
comprising the steps of:
- 25 (a) constructing oligonucleotide molecules that encode peptide epitopes;
(b) annealing the oligonucleotide molecules to form duplexes;
(c) introducing the oligonucleotide duplexes into an expression vector so as
to encode a fusion protein;
(d) introducing the expression vector into a host cell to allow expression of
the fusion protein;
30 (e) isolating the fusion protein produced from a culture of said host cells.
- 29 The method of claim 28, further comprising the step of conjugating the fusion
protein to polysaccharide.

- 30 The method of claim 29, wherein the host cell is an *E. coli* bacterium.







FIG. 2

N6



ATG GGG GGT TCT CAT CAT CAT CAT GGT ATG GCT AGC ATG GAT TAC AAG GAC GAC GAT GAT ATC GAA GGT CGC
Met Gly Ser His His His His His Gly Met Ala Ser Met Asp Tyr Lys Asp Asp Asp Ile Glu Gly Arg

AAA GGT GTT TCC ATC GAC AAA TTC CGT ATC TTC TGC AAA GGT AAC CCG AAA AAA GGT CTG AAA TTC ATC ATC AAA CGT
Lys Gly Val Ser Ile Asp Lys Phe Arg Ile Phe Cys Lys Ala Asn Pro Lys Lys Gly Leu Lys Phe Ile Ile Lys Arg

TAC ACC CCG AAC AAC GAA ATC GAC TCC AAA GGT ATC CGT GAA GAC AAC AAC ATC ACC CTG AAA CTG GAC CGT TGCAAC
Tyr Thr Pro Asn Asn Glu Ile Asp Ser Lys Gly Ile Arg Glu Asp Asn Ile Thr Leu Lys Leu Asp Arg Cys Asn


AAC AAA GGT GAA AAG AAG ATC GCT AAA ATG GAA AAA GCT TCT TCT TCT TCT AAC GTT GTT AAC TCT AAA GGT TTC AAC
Asn Lys Gly Glu Lys Lys Ile Ala Lys Met Glu Lys Ala Ser Ser Val Phe Asn Val Val Asn Ser Lys Gly Phe Asn

AAC TTC ACC GTT TCC TTC TGG CTG CGT GTT CCG AAA GTT TCC GCT TCC CAC CTG GAA AAA GGT CAG TAC ATC AAA GCT
Asn Phe Thr Val Ser Phe Trp Leu Arg Val Pro Lys Val Ser Ala Ser His Leu Glu Lys Gly Gln Tyr Ile Lys Ala

AAC TCC AAA TTC ATC GGT ATC ACC GAA AAA GGT GGA TCC TAA
Gln Ser Lys Phe Ile Gly Ile Thr Glu Lys Gly Gly Ser End

FIG. 2(CONTD.)

N10



ATG GGG GGT TCT CAT CAT CAT CAT GGT ATG GCT AGC ATG GAT TAC AAG GAC GAC GAT GAT ATC GAA GGT CGC
Met Gly Gly Ser His His His His Gly Met Ala Ser Met Asp Tyr Lys Asp Asp Asp Ile Glu Gly Arg

AAA GGT GTT TCC ATC GAC AAA TTC CGT ATC TTC TGC AAA GCT AAC CCG AAA AAA GGT CTG AAA TTC ATC ATC AAA CGT
Lys Gly Val Ser Ile Asp Lys Phe Arg Ile Phe Cys Lys Ala Asn Pro Lys Lys Gly Leu Lys Phe Ile Lys Arg

TAC ACC CCG AAC AAC GAA ATC GAC TCC AAA GGT ATC CGT GAA GAC AAC AAC ATC ACC CTG AAA CTG GAC CGT TGC AAC
Tyr Thr Pro Asn Asn Glu Ile Asp Ser Lys Gly Ile Arg Glu Asp Asn Asn Ile Thr Leu Lys Leu Asp Arg Cys Asn

AAC AAA GGT GAA AAG AAG ATC GCT AAA ATG GAA AAA GCT TCT TCT GTT TTC AAC GTT GTT AAC TCT AAA GGT TTC AAC
Asn Lys Gly Glu Lys Lys Ile Ala Lys Met Glu Lys Ala Ser Ser Val Phe Asn Val Val Asn Ser Lys Gly Phe Asn

AAC TTC ACC GTT TCC TTC TGG CTG CGT GTT CCG AAA GTT TCC GCT TCC CAC CTG GAA AAA GGT CAG TAC ATC AAA GCT
Asn Phe Thr Val Ser Phe Trp Leu Arg Val Pro Lys Val Ser Ala Ser His Leu Glu Lys Gly Gln Tyr Ile Lys Ala

AAC-TCC AAA TTC ATC GGT ATC ACC GAA AAA GGT GGA TCT CCG CAT CAT ACC GCG CTG CCG CAG GCG ATT CTG TGC TGG
Gln Ser Lys Phe Ile Gly Ile Thr Glu Lys Gly Ser Pro His His Thr Ala Leu Arg Gln Ala Ile Leu Cys Trp

GGC GAA CTG ATG ACC CTG GCG AAA GGA TCT CCG AAA TAT GTG AAA CAG AAC ACC CTG AAA CTG GCG ACC AAA GGA TCG
Gly Glu Leu Met Thr Leu Ala Lys Gly Ser Pro Lys Tyr Val Lys Gln Asn Thr Leu Lys Leu Ala Thr Lys Gly Ser

TTT TTT CTG CTG ACC CCG ATT CTG ACC ATT CCG CAG TCT CTG GAT AAA GGC TAT TCT GGC CCG CTG AAA GCG GAA ATT
Phe Phe Leu Leu Thr Arg Ile Leu Thr Ile Pro Gln Ser Leu Asp Lys Gly Tyr Ser Gly Pro Leu Lys Ala Glu Ile

GCG CAG CCG CTG GAA GAT GTG AAA GGA TCC TAA
Ala Gln Arg Leu Glu Asp Val Lys Gly Ser End

FIG. 2(Contd.)



FIG. 4**FIG. 3**

FIG. 5

Coomassie




FIG. 6

Western blot





FIG. 7

FIG. 8